



DRAFT

Great Lakes Binational Strategy for Mercury Risk Management

April 2018

**A document to assist in the engagement of key stakeholders and the public
in strategy development**

This draft was prepared by Environment and Climate Change Canada and
the United States Environmental Protection Agency



Disclaimer

The purpose of this document is to propose mercury risk mitigation and management strategies in accordance with Annex 3 of the Great Lakes Water Quality Agreement (GLWQA). The mention of trade names, commercial products, or organizations does not imply endorsement by the Canadian or United States governments.

Acknowledgments

Funding for this report was provided by the US Environmental Protection Agency (US EPA) by Battelle Memorial Institute under US EPA Contract EP-R5-11-07, Contracting Officer's Technical Representative F. Anscombe, Chicago, IL. This report was prepared with guidance from staff at the US Environmental Protection Agency and Environment and Climate Change Canada.

Cover and Page Banner Photo Credits

Top cover: Peninsula Harbour Remediation, Marathon, Ontario, source: infosuperior.com.

Bottom cover: St. Clair River Area of Concern (AOC) Cleanup, source: Roger Santiago, Environment Canada.

Page banners: Muskegon AOC Coastal Acquisition Project, source: National Oceanic and Atmospheric Administration.

Executive Summary

The Canada-United States Great Lakes Water Quality Agreement (GLWQA) seeks to reduce the anthropogenic release of chemicals of mutual concern (CMCs), including mercury, into the air, water, land, sediment, and biota of the Great Lakes basin ecosystem. Under the GLWQA the Parties have agreed to adopt, as appropriate, the principles of virtual elimination and zero discharge for releases and control of CMCs.

This document provides a Binational Strategy for mercury to focus efforts of the Governments of Canada and the United States, in cooperation and consultation with State and Provincial Governments, Tribal Governments, First Nations, Métis, Municipal Governments, watershed management agencies, other local public agencies, industry, and the public in implementing risk mitigation and management options aimed at reducing mercury in the Great Lakes region. The Parties and their partners will use this strategy as guidance to identify, prioritize, and implement actions to reduce CMCs. Strategy options are organized under five categories: Regulations and Other Risk Mitigation and Management; Compliance Promotion and Enforcement; Pollution Prevention; Monitoring, Surveillance, and Other Research; and Domestic Water Quality. The Parties commit to incorporating, to the extent feasible, options outlined herein in their decisions on programs, funding, and staffing, but implementation would take place by agencies with mandates to undertake work in these areas. As noted in the GLWQA, the Parties' obligations are subject to the appropriation of funds in accordance with their respective procedures.

Mercury is a naturally occurring metallic element that has been used in a variety of applications and is released from both natural and anthropogenic sources. Concentrations of mercury exceed human and ecological risk thresholds in many areas of the Great Lakes. Due to its adverse effects in humans and wildlife, mercury has been widely studied, monitored, regulated, and targeted for action in Canada, the United States, and globally. Yet mercury continues to be a threat to ecosystems and human health.

This strategy identifies multiple gaps that require further development:

- An overall lack of knowledge as to how a changing climate may impact the mercury cycle
- A need to enhance emissions data and apply innovative tools such as isotopic mercury ratios to improve the ability of current models to predict the relationship between mercury emissions and fish concentrations
- A need for comprehensive evaluation of how well existing regulatory programs are working to limit and reduce mercury impacts on the Great Lakes

To address these gaps, this Binational Strategy document proposes multiple options as outlined in **ES Table A**. By implementing these options, stakeholders will be improving the health of the Great Lakes basin and their respective communities.

ES Table A. Summary of the Canada-United States Strategy Options for Mercury

Category of Action				
Regulations and Other Risk Mitigation and Management Actions	Compliance Promotion and Enforcement	Pollution Prevention	Monitoring, Surveillance, and Other Research Efforts	Domestic Water Quality
Strategy Options				
<p>Evaluate the effectiveness of existing regulatory programs to ensure maximum efficiency and overall positive implications on a global scale (Canada)</p> <p>Evaluate the effectiveness of existing emissions regulatory programs for addressing mercury pollution (US)</p> <p>Review and update actions to match current scientific understanding and regional context (Canada and US)</p> <p>Identify manufacturing processes or products that intentionally add mercury (US)</p> <p>Continue to reduce mercury emissions resulting from coal-fired generation of electricity (Canada)</p> <p>Continue implementation of domestic regulations and other risk management activities for mercury (Canada and US)</p> <p>Develop the National Strategy for Safe and Environmentally Sound Disposal of Lamps Containing Mercury (Canada)</p> <p>Continue remediation of mercury-contaminated sites and sediments (Canada and US)</p> <p>Amend the Products Containing Mercury Regulations to further reduce mercury in products (Canada)</p>	<p>Promote compliance with domestic and international mercury activities and initiatives (Canada and US)</p> <p>Continue implementation of respective obligations of the Minamata Convention on mercury (Canada and US)</p>	<p>Enhance public outreach and educate the public and facility staff on potential sources of mercury and proper actions to follow when handling mercury containing products (Canada and US)</p> <p>Enhance public outreach and educate the public on how to obtain and implement site-specific fish consumption advisories (Canada and US)</p> <p>Encourage industries to track their P2 activities and efforts in the National Pollutant Release Inventory (NPRI) or Toxics Release Inventory (TRI), or via P2 promotion activities (fact sheets, case studies) (Canada and US)</p> <p>Highlight pollution prevention successes (Canada and US)</p> <p>Implement best available techniques and best environmental practices for new and substantially modified sources (Canada and US)</p>	<p>Continue monitoring mercury in environmental media in the Great Lakes (air, precipitation, sediment, fish, and other wildlife) and publish results in a variety of publications (e.g., online and open data portals, government reports and scientific journals) to maximize the intended audience (Canada and US)</p> <p>Continue efforts to update and maintain mercury emissions inventories in a manner such that regional and global emissions can be tabulated (Canada and US)</p> <p>Conduct additional research on methylation dynamics and the differential impacts of mercury in nearshore versus offshore environments (US)</p> <p>Enhance existing models to track long-range atmospheric transport and the rate of methylmercury formation in the environment and its corresponding ecological risk (Canada and US)</p> <p>Develop cost-effective, reliable and effective tools (e.g., passive samplers) for collecting long-term mercury multi-media monitoring data (Canada and US)</p> <p>Develop and populate a structured data system to track mercury sources, manifests, waste, and products (Canada and US)</p>	<p>Review and update existing domestic water quality standards, if necessary (Canada and US)</p>

Table of Contents

Disclaimer.....	ii
Acknowledgments.....	ii
Cover and Page Banner Photo Credits	ii
Executive Summary.....	iii
Table of Contents.....	v
Figures.....	vi
Tables	vii
Acronyms and Abbreviations	viii
1 Introduction	1
2 Chemical Profile	2
2.1 Chemical Identity	2
2.2 Physical and Chemical Properties	2
2.3 Environmental Fate and Transport	2
2.4 Sources and Releases of Mercury in the Great Lakes.....	3
2.4.1 Uses and Quantities in Commerce.....	3
2.4.2 Sources of Emissions and Releases.....	4
2.4.3 Mercury in Environmental Media	6
2.4.3.1 In Air	6
2.4.3.2 In Surface Water	6
2.4.3.3 In Sediments	7
2.4.3.4 In Biota	7
2.5 High Level Summary of Risks	8
3 Existing Mercury Management/Control Policies, Regulations, and Programs	9
3.1 United States.....	9
3.1.1 Existing Statutes and Regulations	9
3.1.2 Pollution Prevention Actions	10
3.1.3 Risk Management Actions	11
3.1.4 Monitoring, Surveillance and Other Research Efforts	11
3.1.5 United States Guidelines and Standards	12
3.2 Canada	12
3.2.1 Existing Statutes and Regulations	12
3.2.2 Pollution Prevention Actions	13
3.2.3 Risk Management Actions	13
3.2.4 Monitoring, Surveillance, and Other Research Efforts	14
3.2.5 Canadian Environmental Quality Guidelines and Standards	16
3.3 Binational Actions	18
3.3.1 Great Lakes Binational Toxics Strategy	18
3.3.2 Lakewide Action and Management Plans	18

3.3.3	Great Lakes Regional Collaboration	18
3.3.4	National Atmospheric Deposition Program/Mercury Deposition Network	19
3.3.5	Cooperative Science and Monitoring Initiative	19
3.3.6	State of the Great Lakes Reporting	19
3.4	International	20
3.4.1	United Nations Economic Commission for Europe, Convention on Long-Range Transboundary Air Pollution.....	20
3.4.2	United Nations Environment Program, Minamata Convention on Mercury.....	20
3.4.3	UNEP Global Mercury Partnership.....	21
3.4.4	Arctic Contaminants Action Plan and the Arctic Monitoring and Assessment Program under the Arctic Council.....	21
3.4.5	The Commission for Environmental Cooperation.....	21
4	Gap Analysis	22
4.1	Gaps and Needs for Action	22
4.2	Exceedances of or Non-compliance with Environmental Quality Guidelines	23
5	Risk Mitigation and Management Options to Address Gaps	23
5.1	Regulations, and Other Risk Mitigation and Management	23
5.2	Compliance Promotion and Enforcement	25
5.3	Pollution Prevention	25
5.4	Monitoring, Surveillance, and Other Research Efforts	26
5.5	Domestic Water Quality.....	27
6	Conclusions	27
7	Figures.....	29
8	Tables	34
9	References	39

Figures

Figure 1. The Mercury Cycle. Source: Evers et al. (2011)	29
Figure 2. Decline in Mercury Emissions 1990-2005. Source: Evers et al. (2011).....	29
Figure 3. Trends in National Emissions Inventory (NEI) Mercury emissions (tons) Source: United States EPA 2014 NEI version 2 Technical Support Document (TSD)	30
Figure 4. Historical and Projected Canadian Mercury Air Emissions Trends. Source: ECCC (2016b)	30
Figure 5. Seven-year mean annual mercury wet deposition based on NADP/MDN monitoring data. Source: Evers et al. (2011)	31
Figure 6. Spatial Distribution of Mercury in Great Lakes Sediments. Inset is Lake St. Clair Corridor. Source: State of the Lakes Technical Report (2017)	31
Figure 7. Long-Term Mercury Trends in Chinook Salmon, Coho Salmon, Lake Trout, Lake Whitefish and Walleye in the Great Lakes. Source: State of the Lakes Technical Report 2017.....	32
Figure 8. Great Lakes Fish Consumption Advisories for Mercury. Source: Evers et al. (2011).....	33

Tables

Table 1. Physical and Chemical Properties of Mercury and Select Compounds.	34
Table 2. Estimated Amount of Elemental Mercury in Mercury-Added Products in the United States (2013)	35
Table 3. Sources of Global Mercury Supply, 2015.	35
Table 4. Total Mercury Emissions in Canada by Sector, 1990-2015.	35
Table 5. United States Federal Statutes Authorizing Mercury Regulations.	36
Table 6. Summary of Great Lakes States Mercury Programs from a 2011 Survey.	37
Table 7. United States Standards and Canadian Environmental Quality Guidelines for Mercury and Methylmercury and Average Great Lakes Concentrations.	38

Acronyms and Abbreviations

ACAP	Arctic Contaminants Action Plan
AMAP	Arctic Monitoring and Assessment Program
AMNet	Atmospheric Mercury Network
AOC	Areas of Concern
ASGM	Artisanal and small-scale gold mining
CAMNet	Canadian Atmospheric Mercury Measurement Network
CAPMoN	Canadian Air and Precipitation Monitoring Network
CARA	Clean Air Regulatory Agenda
CCME	Canadian Council of Ministers of the Environment
CEC	Commission for Environmental Cooperation
CEPA	Canadian Environmental Protection Act
CEQG	Canadian Environmental Quality Guidelines
CHMS	Canadian Health Measures Survey
CMC	Chemicals of Mutual Concern
CNR	Consiglio Nazionale delle Ricerche (Institute for Atmospheric Pollution, Italy)
COP1	Minamata Convention on Mercury
CSMI	Cooperative Science and Monitoring Initiative
CWS	Canada-Wide Standards
ECCC	Environment and Climate Change Canada
ECCC-AMM	Environment and Climate Change Canada – Atmospheric Mercury Monitoring
EPCRA	Emergency Planning and Community Right-to-Know Act
FCMSP	Fish Contaminants Monitoring and Surveillance Program
FNBI	First Nations Biomonitoring Initiative
GLBTS	Great Lakes Binational Toxics Strategy
GLEND	Great Lakes Environmental Database
GLNPO	Great Lakes National Program Office
GLRC	Great Lakes Regional Collaboration
GLRI	Great Lakes Restoration Initiative
GLWQA	Great Lakes Water Quality Agreement
Hg	Mercury
HgS	Mercuric Sulfide, Cinnabar
IADN	Integrated Atmospheric Deposition Network
ITT	Identification Task Team
LAMP	Lakewide Action and Management Plan
LRTAP	Long-Range Transboundary Air Pollution
LSBP	Lake Superior Binational Program
MACT	Maximum Achievable Control Technology
MATS	Mercury and Air Toxics Standards
MDN	Mercury Deposition Network
MEBA	Mercury Export Ban Act of 2008
MPO	Manufactured, Processed, or Otherwise Used
NADP	National Atmospheric Deposition Program
NARAP	North American Regional Action Plan
NAtChem	National Atmospheric Chemistry (Database)
NCP	Northern Contaminants Program
Ni-Cd	Nickel Cadmium

NPRI	National Pollutant Release Inventory
NRDC	Natural Resources Defense Council
P2	Pollution Prevention
PCB	Polychlorinated Biphenyl
POTW	Publicly Owned Treatment Works
PRTR	Pollutant Release and Transfer Registry
RCRA	Resource Conservation and Recovery Act
SiGL	Science in the Great Lakes
SOLEC	State of the Lakes Ecosystem Conferences
SSLA	Small Sealed Lead-Acid
TGM	Total Gaseous Mercury
TPM	Total Particulate Mercury
TRI	Toxics Release Inventory
TSCA	Toxic Substances Control Act
UNEP	United Nations Environment Programme
UNIDO	United Nations Industrial Development Organization
US EPA	United States Environmental Protection Agency
US FDA	Food and Drug Administration (US)
USGS	United States Geological Survey
ZDDP	Zero Discharge Demonstration Program

1 Introduction

The purpose of [Annex 3](#) of the Canada-United States Great Lakes Water Quality Agreement (GLWQA) is to reduce the anthropogenic release of chemicals of mutual concern (CMCs) into the Waters of the Great Lakes, recognizing: (1) the importance of life cycle management, (2) that knowledge and information are fundamental to sound management, (3) that CMCs may be managed at the federal, state, provincial, indigenous peoples, and local levels through a combination of regulatory and non-regulatory programs, (4) that international efforts may contribute to reductions from out-of-basin sources, and (5) that the Public can contribute to achieving reductions. While there is no requirement in the GLWQA to set reduction targets, consideration should be given to existing guidelines and the work of other Annexes.

In 2016, the two governments designated mercury as [one of eight](#) CMCs. In designating mercury as a CMC, the Parties have agreed that it poses a threat to the Great Lakes, that current management actions are insufficient, and that further action benefiting the Great Lakes basin is warranted. These actions are documented in binational strategies that may include research, monitoring, surveillance and pollution prevention and control provisions. The purpose of the binational strategies is therefore to reduce releases of CMCs by focusing efforts of Governments, agencies, and the Public in implementing risk mitigation and management actions. The Governments of the United States and Canada are responsible for the implementation of the GLWQA. Within the United States, the US EPA's Great Lakes National Program Office (GLNPO) coordinates these efforts. Within Canada, Environment and Climate Change Canada's Ontario Regional Director General's Office coordinates these efforts.

The Parties and their partners will use this strategy as guidance to identify, prioritize, and implement actions to reduce CMCs. Reductions will only be achievable with widespread on-the-ground action, but it will take time to implement actions to the extent that significant reductions are achieved, and it will take time for the aquatic environment to respond. Factors such as climate change, legacy sources, and changing human activities on the landscape make it difficult to predict the rate at which we could see significant changes in the lakes. The ultimate success of the strategy depends on the combined efforts of the Great Lakes community. The strategy and its implementation will be reviewed on a regular basis and reported through the Progress Report of the Parties. While the GLWQA does not provide timelines for strategy implementation, the strategy should be reviewed periodically. Please note that during the time frame of re-evaluation, no new chemical nominations will be accepted.

This mercury strategy covers a list of 23 management options, in Canada and/or the United States, to address threats to water quality by reducing mercury releases. These options can be used to help identify, support or coordinate ongoing or new projects. The options are organized under five categories: Regulations and Other Risk Mitigation and Management Actions; Compliance Promotion and Enforcement; Pollution Prevention; Monitoring, Surveillance, and Other Research Efforts; and Domestic Water Quality. The Parties commit to incorporating, to the extent feasible, respective options in the CMC strategies in their decisions on programs, funding, and staffing. Implementation will take place, to the extent feasible, by agencies with mandates to undertake work in these areas. As noted in the GLWQA, the Parties' respective obligations are subject to the appropriation of funds in accordance with their respective procedures. Implementation of some CMC actions may be supported through other GLWQA Annexes, for example Annexes 2 (Lakewide Management) and 10 (Science).

2 Chemical Profile

An extensive summary of environmental data and other pertinent information considered part of the process of designating mercury as a CMC is available in the [Binational Summary Report: Mercury](#) produced by the Identification Task Team (ITT) (2015). [The Extent and Effects of Mercury Pollution in the Great Lakes Region, Great Lakes Mercury Connections](#), a synthesis published by the Biodiversity Research Institute with the Great Lakes Commission and the University of Wisconsin-La Crosse, provides an overview of sources, cycling, and impacts of mercury (Evers et al., 2011). The Canadian Mercury Science Assessment Report (ECCC 2016) also provides a comprehensive scientific evaluation and synthesis of mercury in the Canadian environment.

2.1 Chemical Identity

Mercury (Hg) is a naturally occurring metallic element. There are three forms of mercury: elemental, inorganic, and organic mercury compounds. Elemental mercury is a heavy, silvery-white liquid metal, and is the only metallic element that is liquid at room temperature. Inorganic mercury compounds occur when mercury combines with elements other than carbon, such as chlorine, oxygen, or sulfur. Inorganic mercury compounds, also called mercury salts, are mostly white powders or crystals, except for mercuric sulfide (cinnabar), which is red and turns black after exposure to light. Mercury combined with carbon forms organic mercury compounds; methylmercury is the most common organic mercury in the environment (US EPA, 2017b).

2.2 Physical and Chemical Properties

Mercury is valued for use in industrial applications due to its physio-chemical properties of high surface tension, high specific gravity, low electrical resistance, high reflectance, and constant volume of expansion in the liquid state (Schroeder and Munthe, 1998; Park and Zheng, 2012). Compared with other metals, mercury is a poor conductor of heat and a fair conductor of electricity. Select physical and chemical properties of mercury and common mercury compounds are shown in **Table 1**. When naturally occurring, mercury is a metal that is principally in the form of the inorganic mineral cinnabar (HgS) (Michigan DEQ and US EPA, 2013). The cinnabar form is an insoluble and stable compound (ATSDR, 2013). However, natural or anthropogenic releases of inorganic mercury to the environment allow for natural biological processes to convert inorganic mercury to the more toxic organic form of methylmercury (Schroeder and Munthe, 1998).

2.3 Environmental Fate and Transport

The simplified mercury cycle illustrated in **Figure 1** shows how mercury enters and cycles through ecosystems, biomagnifies in the food chain, and bioaccumulates in fish and wildlife (Evers et al., 2011). Within the aquatic environment, inorganic mercury can be converted into methylmercury through a series of complex processes often involving sulfate-reducing bacteria typically in wetlands and sediments, (Evers et al., 2011). Methylmercury bioaccumulates and biomagnifies through the food chain, with levels increasing from small aquatic organisms, to small fish to larger fish, to wildlife. This may create a significant source of dietary exposure to humans who consume fish or wildlife. Top predatory fish such as walleye and lake trout can have mercury concentrations more than one million times higher than that of the surrounding water (International Joint Commission, 2015).

Atmospheric emissions of mercury can result from natural processes, including forest fires and volcanic activity, and anthropogenic processes such as the combustion of coal or industrial activities (Cohen et al., 2007; Evers et al., 2011). Mercury emitted from anthropogenic sources may remain in the atmosphere for six months to a year, enabling long-range transport (International Joint Commission,

2015) prior to eventual atmospheric deposition (Evers et al., 2011). Deposition can take the form of wet or dry deposition. In wet deposition, mercury is removed from the atmosphere and deposited back on the Earth in the form of precipitation (e.g., rain, snow) (Evers et al., 2011). In dry deposition, mercury is removed from the atmosphere and deposited as a gas or particle (Evers et al., 2011). As a result of wet or dry deposition, mercury can accumulate on trees, soil, water, or other surfaces (Cohen et al., 2007). In addition to long-range transport and deposition, mercury can also deposit locally, depending primarily on its oxidation state. Gaseous oxidized and particulate mercury forms generally deposit much more rapidly than elemental mercury and have a much shorter atmospheric residence time (on the order of days to weeks). Although the oxidized forms make up a small fraction of total atmospheric mercury, they can be a large part of total mercury deposition (wet +dry).

2.4 Sources and Releases of Mercury in the Great Lakes

Natural emissions of mercury result from its presence in the Earth's crust and are produced by volcanic and geothermal activity, soils enriched with mercury-containing minerals, forest fires, and erosion of mineral deposits.

Atmospheric emission and deposition is now identified as the greatest source of mercury to the Great Lakes region (Evers et al., 2011). Atmospheric mercury can travel from distant global and regional sources, or can be released locally and remain in the Great Lakes region (Evers et al., 2011). An isotopic mixing model (Lepak et al. 2015) identified three primary mercury sources for sediments: atmospheric, industrial, and watershed-derived. Results indicate atmospheric sources dominate in Lakes Huron, Superior, and Michigan sediments while watershed-derived and industrial sources dominate in Lakes Erie and Ontario sediments. Comparison of signatures in predatory fish from three lakes reveals that bioaccumulated mercury is more isotopically similar to atmospherically derived mercury than a lake's sediment. An improved global emission inventory for the period 1990 to 2010 (Zhang et al. 2016) found a 20% decrease in total mercury emissions and a 30% decrease in anthropogenic Hg₀ emissions, with much larger decreases in North America and Europe offsetting the effect of increasing emissions in Asia. The large trends observed in North America and Europe reflect the phase-out of mercury from commercial products as well as the co-benefit from SO₂ and NO_x emission controls on coal-fired utilities.

2.4.1 Uses and Quantities in Commerce

Historically, mercury-added products have been widely used in residential, commercial, industrial, military, marine, and medical environments. However, since 1980, the use of mercury in products sold in the United States has decreased more than 97 percent. Major remaining mercury-added product categories for the United States are listed in **Table 2** (US EPA 2017c, IMERC). These include industrial products and components, such as switches and relays, dental amalgam, thermostats, lamps, button cell batteries, and formulated products such as coating materials, acids, alkalis, bleach, pharmaceutical products, stains, reagents, preservatives, cosmetics, and dyes (Carpenter et al., 2011). Current global uses for commodity mercury include vinyl chloride monomer (VCM) production, small-scale artisanal gold mining (ASGM) operations, mercury cell chlor-alkali processes, and various consumer goods including compact fluorescent light bulbs and other lamps (Carpenter et al., 2011). The US does not use mercury VCM or ASGM processes and Canada does not engage in either process.

In Canada, the *Products Containing Mercury Regulations* prohibit the import and manufacture, since November 8, 2015, of products containing mercury, with some exemptions for essentials products that have no technically or economically viable alternatives. Mercury and its compounds are listed as prohibited ingredients on the [Cosmetic Ingredient Hotlist](#). As such, their use is not permitted in cosmetic products in Canada. Data received during the 2017 reporting period indicate that products containing

mercury imported or manufactured in Canada during 2016 represented approximately one metric tonne of mercury and were mainly dental amalgam.

The only known current use of elemental mercury in a U.S. manufacturing process is in the production of chlorine and caustic soda using mercury cell technology. Most U.S. facilities that used the mercury cell process have closed or converted to mercury-free manufacturing technologies; the number of operating mercury cell plants in the United States has declined from 14 facilities in 1996 to 2 facilities in 2013. An estimated 368 metric tons of elemental mercury were in use by the chlor-alkali industry in 2013 (U.S. EPA 2017c). There are no mercury cell chlor-alkali plants in Canada; the last mercury cell chlor-alkali plant closed in 2008 (ECCC, 2016a).

In 2017, UNEP estimated the average global supply of mercury in 2015 at approximately 3480-4785 metric tons per year (UNEP 2017). Mining of metallic mercury contributes to the global supply of mercury. Mercury is currently mined only in China, Mexico, Indonesia and the Kyrgyz Republic (European Commission, 2017). The main sources of mercury for the global market, based on 2007 data, are summarized in **Table 3**.

The industrial use of mercury is increasingly recognized as posing a liability, therefore the United States and other countries have begun to accumulate a large surplus of commodity mercury (i.e., manufactured or recovered and sold or stored for later use)¹ (Carpenter et al., 2011). According to US EPA (2017), there are currently two primary sources of commodity mercury supply in the United States:

- Byproduct from metal mining and processing
- Recovery from treatment of waste

US EPA estimates byproduct elemental mercury production at 12 metric tons in 2011 and the quantity of mercury recovered from treatment of hazardous waste to be 66 metric tons in 2013 (US EPA 2017c).

According to the Associations of Lighting and Mercury Recyclers (ALMR), more than 60 companies recycle mercury-containing products in the United States. There are facilities in the Great Lakes Region in Wisconsin, Minnesota, Illinois and Pennsylvania however not all of these facilities retort mercury, but only recover mercury for further processing. Four main recycling companies specialize in retorting or reprocessing mercury waste to commodity-grade mercury. One of these is located in Illinois; two are located in Pennsylvania and one in Minnesota (US EPA 2007).

There are no gold, silver or zinc mining operations in the Great Lakes basin. Mercury from the iron mining industry (stack emissions from taconite processing) in northeastern Minnesota is the principle mining related source in the Great Lakes basin. According to Berndt (2003), mercury emissions in 2000 of 342 kg/year were predominantly elemental and hence not deposited locally. According to the Iron Mining Association of Minnesota there are currently eight companies mining or planning to mine iron ore in Minnesota (<http://taconite.org/mining-industry/mines>).

2.4.2 Sources of Emissions and Releases

Both natural and anthropogenic activities contribute to the mercury cycle. Natural sources of mercury, such as volcanic eruptions and emissions from the ocean, have been estimated to contribute about one-

¹ Commodity mercury is defined as 99.99% pure by volume Carpenter, C., L. O'Connor, J. Elmer and D. DePinho (2011). Assessing the Impacts of the Mercury Export Ban Act of 2008 on the U.S. Mercury Recycling Industry. WM11 Global Achievements and Challenges in Waste Management, Phoenix, AZ.

third of current worldwide mercury air emissions, whereas anthropogenic emissions account for the remaining two-thirds (GLRC, 2010). Anthropogenic emissions, once deposited, can be re-emitted. Like natural emissions, these re-emissions are believed to be largely in the form of elemental mercury (Cohen et al., 2007). Much of the mercury circulating through the environment today is likely mercury that was released years ago when mercury was frequently used in many industrial, commercial, and residential products and processes (GLRC, 2010) although the ocean has become a significant sink (Lamborg et al. 2014).

Initial regulatory attention to sources of mercury in the 1970s focused on large industrial chlor-alkali plants and pulp and paper mills that discharged mercury directly to the Great Lakes and to the rivers and streams draining into the lakes. Mercury has also been emitted into the atmosphere by municipal and medical waste incinerators (NYSDEC, 2017). Atmospheric emissions and deposition were identified as the largest source of mercury to the Great Lakes region in a 2010 report (GLRC, 2010), with the greatest proportion of mercury released into the atmosphere originating from coal-fired electric power plants. Coal-fired electric generating plants were phased out in Ontario in 2014. However, coal-fired electric generating plants are currently in use in states in the Great Lakes basin.

As shown in **Figure 2**, there was a 50% decline in anthropogenic mercury emissions in the Great Lakes states between 1990 and 2005 (Evers et al., 2011). This decline is likely the result of provincial, state, regional, binational, and voluntary actions to control mercury emissions. Canadian inventories reported an 85% decrease in mercury air emissions between 1990 and 2010 (**Table 4**) (ECCC, 2016a). As of 2011, coal-fired power plants were identified as the largest source of anthropogenic mercury atmospheric emissions in the Great Lake States, contributing approximately 39% of total anthropogenic mercury emissions (NEI, 2011). Coal-fired electricity generating plants in Canada are in the process of being phased out (Sibbald, 2016) and were closed in Ontario by 2014. The Mercury and Air Toxic Standards (MATS) and other factors (e.g., fuel switching to natural gas) have contributed to a significant decline in mercury emissions from coal-fired power plants in the US. Overall, mercury emissions sources have decreased in Canada, the United States, and Europe (Evers et al., 2011). However, global emissions of mercury are on the rise, mainly due to contributions from Asia (Evers et al., 2011). It has been estimated that mercury emissions from Asia now account for ~50% of the total anthropogenic emissions (ECCC, 2016a).

In 2005, Metal ore processing was the second largest contributor and accounted for 14% of atmospheric mercury emissions in the Great Lakes states (ITT, 2015). Coal-fired plants were identified as the highest individual contributors at the state or provincial level in the Great Lake region, with the exception of Minnesota and New York (Evers et al., 2011), although since closing its remaining coal fired plants in 2014, Ontario is no longer contributing to this source. Other stationary sources are medical and municipal waste incinerators and industrial boilers (Evers et al., 2011). Other major anthropogenic sources of mercury include the use and disposal of mercury-containing products, processing of metals, and the manufacturing of cement (Michigan DEQ and US EPA, 2013).

By using stable mercury isotopes, researchers are now able to assess sources of mercury in Great Lakes sediments. Mercury isotopes in Great Lakes precipitation can also be used as a tool for source attribution (Sherman and Blum, et al., 2016). Atmospheric (precipitation)-derived mercury sources dominate the sediments of Lakes Superior, Huron, and Michigan, whereas watershed and industrial mercury sources dominate the sediments of Lakes Erie and Ontario (Evers et al., 2011). Mercury can also be introduced to water bodies from water treatment facilities or landfill leachate that becomes contaminated from the use and disposal of consumer products containing mercury, such as batteries,

light bulbs (i.e., fluorescent lights and compact fluorescent bulbs), and electrical switches (US EPA, 2017b). Much of the point-source mercury pollution from chlor-alkali facilities has been controlled, allowing partial recovery of the Great Lakes, as shown by lower mercury concentrations in lake sediments in the lower Great Lakes (e.g., Lake Ontario), declines in fish mercury concentrations since the 1970s, and the advent of regulatory attention (Evers et al., 2011).

2.4.3 Mercury in Environmental Media

Mercury monitoring in the Great Lakes basin has been ongoing; highlights of existing data are summarized below. Mercury has been detected in various environmental media, with the highest concentrations of mercury being found near urban and industrial areas.

2.4.3.1 In Air

Ambient mercury concentrations in the Great Lakes have decreased significantly since the 1970s (ITT, 2015). Atmospheric mercury concentrations have decreased about 2% per year since 2005 as measured in Canada's Experimental Lakes Area (west of Lake Superior). Wet deposition measurements from the North American Mercury Deposition Network follow these trends with fluxes decreasing about 1.6% per year since 1996 (ECCC and US EPA, 2017). Mercury emissions in the United States have decreased from an estimated 246 tons in 1990 to 52 tons in 2014 (US EPA, 2014) (**Figure 3**). In Canada, emissions have decreased from approximately 35,000 kg in 1990 to approximately 4,400 kg in 2015 (**Figure 4**) (ECCC, 2016b). Reductions in Canada are mostly due to one facility changing from pyrometallurgical to hydrometallurgical zinc production and, to a smaller extent, to increased emission control measures, improved PM emission controls and fuel switching (ECCC, 2018a).

A recent review of mercury in air reported by Cole et al. (2014) described long-term trends in mercury in air and precipitation in Canada. Declines were noted in total gaseous mercury and mercury in precipitation from the 1990s to early 2010 (Cole et al., 2014). Total gaseous mercury concentrations, with some identified exceptions, were generally 1.2 to 1.9 ng/m³ throughout Canada. Wet deposition amounts averaged between 0.1 and 0.8 µg/m² month (Cole et al., 2014).

2.4.3.2 In Surface Water

Although reported present in trace amounts in Great Lakes surface waters, mercury remains a concern due to the potential for bioaccumulation and biomagnification in the food chain (ITT, 2015). Between 2003 and 2009, mercury concentrations in offshore surface waters of the Great Lakes were relatively low. Reported mercury concentrations ranged from 0.3 ng/L to 0.54 ng/L, significantly below the Canadian Water Quality Guidelines for Protection of Aquatic Life value of 26 ng/L (ITT, 2015). The lowest concentrations were identified in Lake Huron and Georgian Bay at 0.24 and 0.3 ng/L, respectively. Lake Superior and Lake Ontario concentrations were reported to be 0.35 ng/L, and Lake Michigan and Lake Erie had the highest concentrations at 0.49 ng/L and 0.54 ng/L, respectively (ITT, 2015).

Studies have identified a large decline (50-75%) in total aqueous mercury concentrations across the Great Lakes since 2000 (ITT, 2015). The significant decline in mercury concentrations has been attributed to concurrent declines in atmospheric deposition, reduced tributary loadings, and increased volatilization of gaseous mercury. However, the rate of decline in mercury concentration in near-shore (<100 meters in depth) waters is outpacing the rate of decline in mercury concentration in off-shore waters. This discrepancy has been identified for all Great Lakes, except for Lake Huron (ITT, 2015).

2.4.3.3 In Sediments

Sediment core samples from Lake Michigan suggest mercury influx peaked around the mid-1900s, then decreased after the 1970s, coinciding with establishment of the Clean Water Act. Evers et al. (2011) noted that the spike and eventual decline in mercury concentrations is consistent with trends in mercury emission and deposition in the Great Lakes. The consistent region-wide decline in mercury sediment levels suggests that local, regional, and binational emission controls have been effective in decreasing the supply of mercury to the Great Lakes basin. The source contribution of mercury to sediments from wet deposition is shown in **Figure 5**.

Between 1970 and 2010, concentrations of mercury in surface sediments in open-lake areas of the Great Lakes ranged from 0.1 to > 1.0 µg/g (dry weight), with the highest concentrations found in Lakes Erie and Ontario, likely due to the influence of industrial activities (ITT, 2015). The spatial distribution of mercury in Great Lakes sediments is shown in **Figure 6**.

Mean mercury concentrations in Lake Erie and Lake Ontario sediment exceeded the Canadian Sediment Quality Guidelines for Protection of Aquatic Life probable effects level (0.486 µg/g dry weight) and threshold effects level (0.17 µg/g dry weight) for study data reported in 2003, 2005, and 2008 (ITT, 2015). In Lake Superior and the Georgian Bay, the mean mercury sediment concentration only exceeded the threshold effects level. In Lake Huron, the mean mercury sediment concentration was below both the probable effects level and threshold effects level (ITT, 2015). The higher mercury concentrations reported in the eastern Great Lakes is likely attributable to greater contributions from current local sources (e.g., electrical power generation) and higher deposition rates near the release locations (ITT, 2015).

2.4.3.4 In Biota

Monitoring efforts over the last forty years by ECCC showed dramatic declines in mercury levels in herring gull (*Larus argentatus*) eggs and multiple fish species (lake trout [*Salvelinus namaycush*], walleye [*Sander vitreus*], rainbow smelt [*Osmerus mordax*]) during the first three decades of monitoring. In the 2000s, the rate of mercury decline slowed, stopped, or even reversed for some monitored sites and species (**Figure 7**). However, there has been an upward trend in mercury levels in two of the top predatory fish species, walleye and lake trout (ITT, 2015). Concentrations of mercury in top predatory fish measured by ECCC between 2008 and 2012 ranged from 233 ng/g wet weight in Lake Superior to 121 ng/g wet weight in Lake Erie (McGoldrick and Murphy, 2016). Concentrations recently observed and reported through U.S. Great Lakes Fish Monitoring Program are consistent, with concentrations of mercury generally between 110 to 250ng/g across all lakes, except for Lake Superior, where elevated concentrations were observed, up to 415ng/g (Carlson and Swackhamer, 2006; US EPA, 2014a).

It should be noted that over 97% of these recently observed sport fillet mercury concentrations are below the 0.5µg/g (500ng/g) target established under the 1987 Canada-United States Great Lakes Water Quality Agreement (Bhavsar et al., 2010; Zanaski et al., 2011; US EPA, 2014a). While it is evident that a large number of consumption advisories still exist on the basis of mercury concentrations, these are generally only minimally (i.e. 8 meals / month) to moderately restrictive (4 meals / month) (US EPA, 2014a).

Herring gull eggs from the Great Lakes have been monitored for persistent toxic chemicals (ITT, 2015). From an egg survey conducted in 2009 across 15 sites in the Great Lakes region, mercury concentrations ranged from 0.0064 µg/g wet weight at Chantry Island, Lake Huron to 0.246 µg/g wet weight at Middle Island, Lake Erie (ITT, 2015). However, the data indicate a decline in mercury concentrations at 14 of the

15 sites over 35 years (1974-2009) that ranged from 22.6 to 85.8%, with one site in Lake Erie showing an increase of 10.5% (ITT, 2015).

Based on analysis of data from 1999 to 2009, Zanaski et al. (2011) determined that average fish total mercury concentrations over this time period in the five lakes were significantly different from one another with the greatest concentrations in Lake Superior, then Huron, Michigan, Ontario and Erie. Evaluation of data from the mid-1970s to 2007 showed that concentrations of mercury generally declined over these three decades but that in recent years, concentrations across the Great Lakes have leveled off and increased slightly in Lake Erie (Bhavsar et al. 2010, International Joint Commission, 2015). A more recent analysis of combined Great Lakes data (except Lake Erie) showed a significant decreasing trend in the lake trout mercury concentrations between 2004 and 2015 with an annual decrease of 4.1% per year, consistent with the decline in regional atmospheric mercury emissions and water mercury concentrations (Zhou et al. 2017). Their analysis detected a breakpoint with a significant decreasing slope (-8.1% per year) before the breakpoint (2010), and no trend after the breakpoint. Examination of individual lakes showed that Lakes Superior and Huron, which are dominated by atmospheric mercury inputs and are more likely than the lower lakes to respond to declining emissions from areas surrounding the Great Lakes, have significant decreasing trends with rates between 5.2 and 7.8% per year from 2004 to 2015. These declining trends appear to be driven by decreasing regional atmospheric Mercury emissions although they may be partly counterbalanced by other factors, including increasing local emissions, food web changes, eutrophication, and responses to global climate change. Lakes Michigan, Erie and Ontario may have been more impacted by these other factors and their trends changed from decreasing to non-decreasing or increasing in recent years.

Since 1993, levels of mercury in walleye from the west basin of Lake Erie have been increasing at 3.4% per year (SOLEC, 2017). Recent studies of lake trout mercury concentrations in Lake Superior showed mercury levels in fish and other Great Lakes biota to be declining at a rate of approximately 6-7% per year since 2003 (Jeremiason, 2017). While there are currently no binational targets for mercury in fish, the observed concentrations are generally below levels of concern for the health of fish-consuming wildlife (500 ng/g wet weight) as established in the 1987 GLWQA (McGoldrick and Murphy, 2016). In Ontario, there is evidence of increasing mercury levels in fish based on analysis of tissue data for the period 2000 to 2012 but these results are predominantly for inland lakes in northern Ontario and shield lakes in southern Ontario (Ghandi et al. 2015).

2.5 High Level Summary of Risks

Once in the environment, mercury can convert to its highly toxic and bioavailable organic form, methylmercury. Methylmercury is an ecological risk to benthic (sediment-dwelling) organisms, fish, birds, and mammals because it bioaccumulates and biomagnifies as it moves up through the food chain. Adverse effects on wildlife include impaired development of early life stages, slower growth, endocrine and reproductive failures, and lethality. There is also evidence that methyl mercury may exhibit similar neurotoxicity endpoints in some species of wildlife as in humans (Basu and Head, 2010).

Methylmercury is a human health risk because it is an easily absorbed and distributed neurotoxin that affects the central nervous system. Consumption of fish and other seafood is the primary pathway for human exposure to methylmercury. Elevated mercury concentrations are the cause for some fish consumption advisories in all the Great Lakes. In Ontario's Great Lakes waters, mercury accounts for 25% of advisories for Lake Superior, 21% for Lake Huron, 40% for Lake St. Clair and the St. Clair River, 11% for Lake Erie, and 12% for Lake Ontario (**Figure 8**) (Evers et al., 2011). Mercury concentrations in the selected sizes of fish from the Great Lakes are now below 0.2 – 0.3µg/g which allows for consumption

advisories that are minimally (i.e., 8 meals per month) to moderately restrictive (i.e., 4 meals per month) (ITT, 2015). Updated fish consumption advisories are available on-line (Great Lakes Consortium for Fish Consumption Advisories, 2018; MOECC, 2018).

Mercury-induced toxicity affects the kidneys and central nervous system. At high-exposure levels, respiratory, cardiovascular, and gastrointestinal effects also occur (ASTDR, 1999; ATSDR, 2013). In pregnant women, methylmercury can cross the placenta into the fetus and accumulate in the fetal brain and other tissues. It can be passed through breast milk from the mother to infants and young children, who are especially vulnerable as their nervous systems are still developing (ASTDR, 1999).

Population-based exposure surveys in the United States have estimated that ~6-8% of childbearing aged women (16-49 years old) are at risk due to elevated methylmercury in their diets (Mahaffey, 2005). Despite mercury concerns, eating fish (especially fish low in mercury) is still recommended as part of a healthy diet, and breast-feeding is still considered the healthiest way to feed an infant (US FDA 2014; Ginsberg 2016; Raymond et al., 2017). “[Eating Well with Canada’s Food Guide](#)” recommends the consumption of at least two, 75 gram, servings of fish each week, and choosing fish that are low in mercury and high in beneficial fatty acids, such as char, herring, mackerel, salmon, sardines and trout.

3 Existing Mercury Management/Control Policies, Regulations, and Programs

Working in partnership, the United States, Canada, the European Union, and individual countries have established strategies to significantly reduce mercury in mercury-containing products, promoted public awareness of adverse health and environmental effects, and proposed legislation to improve management of mercury use and waste.

3.1 United States

3.1.1 Existing Statutes and Regulations

A multitude of regulations have been established at the federal, state, and local levels to limit the availability, use, discharge, emissions and overall number of mercury sources in the United States. In addition to federal mandates, some of the Great Lakes states, Tribes, and local governments (e.g., publicly owned treatment works [POTWs]) have issued more stringent regulations and processes to limit mercury releases.

Notable federal laws that authorize regulations for mercury in the United States include: the Mercury Export Ban Act of 2008 (MEBA), the Mercury-Containing and Rechargeable Battery Management Act, the Toxic Substances Control Act (TSCA), the Clean Air Act, the Clean Water Act, the Emergency Planning and Community Right-to-Know Act (EPCRA), the Resource Conservation and Recovery Act (RCRA), the Safe Drinking Water Act, and the Comprehensive Environmental Response, Compensation and Liability Act (or “Superfund”). In July 2017, the Effluent Limitations Guidelines and Standards for the Dental Category also came into effect. **Table 5** briefly summarizes these federal statutes as they pertain to mercury; additional information can be found at the provided websites.

Prior to 2016, the TSCA provided US EPA with authority to require reporting, recordkeeping and testing requirements, and restrictions relating to chemical substances and/or mixtures. On June 22, 2016, the Frank R. Lautenberg Chemical Safety for the 21st Century Act (Pub. L. 114-182, 130 Stat. 448; Lautenberg Act), amended the TSCA. The updated Act specified that consistent funding be available to enable the US EPA to evaluate and enforce chemical risk requirements, and increase public transparency and education. An additional three objectives in the Act were specific to mercury and mercury compounds:

- **The ban on exporting elemental mercury, previously initiated under MEBA, now includes mercury compounds.** The US EPA published a notice of this statutory prohibition in the Federal Register in August 2016: <https://www.regulations.gov/docket?D=EPA-HQ-OPPT-2016-0411>.
- **Provisions on long-term storage of elemental mercury were amended.** The deadline for the U.S. Department of Energy to open a facility for storage of elemental mercury was extended to 2020.
- **US EPA is required to publish an inventory of mercury supply, use, and trade in the United States and update this inventory every three years.** The US EPA published its initial inventory report in the Federal Register in March 2017: <https://www.regulations.gov/docket?D=EPA-HQ-OPPT-2016-0411>.
- **US EPA is directed to publish a reporting rule to assist in the preparation of the mercury inventory.** The rule, which must be published in June 2018, will require manufacturers to report on their use of mercury and mercury compounds. <https://www.regulations.gov/document?D=EPA-HQOPPT-2017-0421-0001>.

The United States has also made progress on regulatory efforts to decrease mercury emissions from source categories. Under the Clean Air Act, the US EPA established a rule known as the Mercury and Air Toxics Standards (MATS), which includes technology-based emissions limitation standards based on Maximum Achievable Control Technology (MACT) for mercury and other toxic air pollutants from power plants. MATS is estimated to prevent about 90 percent of the mercury in coal burned in power plants from being emitted to the air (US EPA, 2017d). US EPA also has established MACT standards to control mercury emissions from other industry sectors, including municipal waste combustors, medical waste incinerators, gold ore processing plants, Portland cement production facilities, sewage sludge incinerators, electric arc furnace steelmaking facilities, mercury cell chlor-alkali facilities, and industrial boilers. Both new and existing sources are affected. Since the primary source of mercury to the lakes is atmospheric deposition that is either directly deposited to the lake or flows into the lakes through run-off/stream flow, these actions that address mercury emissions in the United States help considerably toward reducing and preventing mercury loadings to these waterbodies.

3.1.2 Pollution Prevention Actions

The US EPA is responsible for implementing the Pollution Prevention (P2) Act of 1990, which involves reducing or eliminating waste at the source by modifying production processes, promoting the use of nontoxic or less toxic substances, implementing conservation techniques, and reusing materials rather than putting them into the waste stream. In many cases, individual states have been leaders in banning mercury in products, in organizing mercury collections, and in working with small businesses to improve handling of mercury. **Table 6** presents findings from a 2011 survey of mercury activities by individual Great Lakes State agencies. Each of the Great Lakes states has additional mercury programs, beyond federal regulations. Each state has one or more programs to assist in the management of mercury-containing products. In addition, all states participate in the National Vehicle Mercury Switch Recovery Program and individually disseminate Fish Consumption Advisories to their citizens (Quicksilver Caucus, 2012).

The US EPA is charged under the P2 Act to integrate P2 policy into its environmental programs, including air, water, toxics, and hazardous waste, as well as to promote P2 practices and source reduction

approaches at other federal agencies. The US EPA established the National Vehicle Mercury Switch Recovery Program, a partnership agreement with the “Big Three” auto manufacturers (i.e., Ford, GM and Chrysler) and junkyards/recyclers, scrapyard shredders, steel mills, and states, which provides for the removal of mercury switches from junked cars before they are shredded and sold as scrap metal to steel mills. Since 2006, the national program has collected more than 4.5 million switches. In the Great Lakes states, nearly 2.3 million switches have been recovered, equating to 9,225 pounds of collected mercury (US Ecology, 2017).

The US EPA's Pollution Prevention Strategic Plan identifies opportunities for the P2 program to help reduce emissions of greenhouse gases, the use of hazardous materials, and the use of natural resources while working toward a greener and more sustainable economy. The P2 Program Strategy for Electronics addresses the problem of environmental impacts of electronics products, from manufacturing processes through use to end-of-life (recycling), all factors that require the use of energy and may result in significant greenhouse gas and other emissions, including mercury from coal-fired energy generation (US EPA, 2010).

3.1.3 Risk Management Actions

The Great Lakes Restoration Initiative (GLRI) is a United States initiative launched in 2010 to provide funds for accelerating efforts to protect and restore the Great Lakes ecosystem. The GLRI is a federal agency coordination of the Great Lakes Interagency Task Force and the Great Lakes Regional Working Group, led by the US EPA. One of the five major areas of the GLRI is Toxic Substances within Areas of Concern (AOCs), and mercury is one of the toxic substances under this focus area. Work accomplished during the GLRI's first five years is described in the Report to Congress and the President (Great Lakes Interagency Task Force, 2015). Between 2004 and 2015, an estimated 4 million cubic yards (3 million cubic meters) of contaminated sediments in United States AOCs were remediated. The remediated sediments contained mercury and other contaminants of concern (US EPA, 2016a). As a result of remediation efforts during GLRI, three AOCs were removed from the list of areas designated as the most contaminated sites on the Great Lakes.

3.1.4 Monitoring, Surveillance and Other Research Efforts

Environmental monitoring and surveillance of Great Lakes have been conducted through several United States parties and coordinated with Canadian program activities to ensure basin-wide coverage and to minimize duplication of efforts. Local, regional, institutional, Tribal, and federal entities have conducted independent and cooperative studies assessing the conditions and status of the Great Lakes for many years. In addition, GLNPO supports work on toxic chemicals, including mercury, with other partners via grants, interagency agreements, and collaborations to address chemical issues as they relate to human health. The following are highlighted programs being done to acquire mercury data within the United States:

- The Great Lakes Open Lakes Trend Monitoring Program began as a component of the [Great Lakes Fish Monitoring and Surveillance Program](#). The program has monitored contaminant trends in fish of the Great Lakes since its inception in the late 1970s. There are 10 permanent monitoring stations in the program. Fish samples are collected at five stations per year, one in each Great Lake. The program aims to collect 50 top predators to be analyzed as 10 five- fish composites, limited to a strict size range, per station each year. Total mercury is one of the chemical parameters analyzed in the fish tissue.
- National Atmospheric Deposition Program/Atmospheric Mercury Network (AMNet) operates 20 sites in the United States, with three AMNet sites located in the Great Lakes basin, which

monitor the atmospheric concentrations of speciated mercury fractions, gaseous elemental mercury, gaseous oxidized mercury, and particulate-bound mercury. These data have supported dry deposition estimates, emission regulatory assessments, model evaluation, and long-term trends (NADP, 2011).

The reporting of results from GLNPO programs and projects is conducted through a variety of mechanisms, including the State of the Lakes Reports, LAMPs, and peer-reviewed literature. Mercury monitoring data from various programs are tracked through three primary databases: Great Lakes Environmental Database (GLENDA), Science in the Great Lakes (SiGL) Mapper, and the Toxics Release Inventory (TRI) Program. The following monitoring networks, programs, and data repositories include collection of mercury data in the Great Lakes region:

- Much of the United States data collected has been placed in GLENDA, which collects and stores environmental data maintained by GLNPO. Air, water, biota, and sediment data are all compiled in the system for users of Great Lakes data (US EPA, 2016b).
- [SiGL Mapper](#) is an additional searchable meta database tool developed by the US Geological Survey (USGS) that allows Great Lakes stakeholders to coordinate and collaborate monitoring and restoration activities on the Great Lakes (US EPA, 2015b). These databases enable researchers to use historic data from across the region to solve complex chemical, biological, and physical relationships that might lead to more advanced methods for pollution identification and remediation actions.
- The [TRI](#) database is the United States' Pollutant Release and Transfer Registry (PRTR) (US EPA, 2014b). The TRI database tracks releases, disposals, and transfers of mercury (and its compounds) from industrial sources that exceed applicable thresholds. Under EPCRA of 1986 all manufacturing or processing facilities that use mercury are required to report the annual quantities of hazardous or toxic chemicals used, as well as pollution prevention and recycling information to the TRI database. This database, maintained by the US EPA, is publicly available and intended to inform the public and help increase the public's knowledge of and access to information on chemicals at individual facilities, their uses, and releases into the environment. States and communities, working with facilities, can use the information to improve chemical safety and protect public health and the environment (US EPA, 2014; US EPA, 2015a).

3.1.5 United States Guidelines and Standards

Table 7 lists US EPA and United States Food and Drug Administration (US FDA) standards and recommendations in effect for mercury in blood, environmental waters, drinking water, groundwater, and fish. There are no ambient air quality recommendations.

3.2 Canada

3.2.1 Existing Statutes and Regulations

In February 2017, Canada introduced comprehensive restrictions on the exports of mercury. The *Export of Substances on the Export Control List Regulations*, under CEPA 1999, restrict the export of mixtures containing elemental mercury at a concentration of 95% or more by weight, with some exemptions, which are in line with the Minamata Convention.

On February 11, 2017 and pursuant to subsection 54(4) of CEPA1999, the Minister of Environment and Climate Change published a *Notice of the Code of Practice for the Environmentally Sound Management of End-of-life Lamps Containing Mercury* in the *Canada Gazette*, Part 1 (ECCC, 2017b). The Code of Practice is designed to encourage collectors, transporters, and recyclers to incorporate best practices in their management of end-of-life mercury-containing lamps to prevent releases of mercury to the environment. Recognizing that northern and remote regions often face unique challenges that can make it difficult to collect and manage end-of-life mercury-containing lamps, the Code includes additional information on management options that can be used to facilitate the implementation of best practices.

The *Products Containing Mercury Regulations*, which came into force in 2015 prohibit the manufacture and import of products containing mercury or any of its compounds, with some exemptions for essential products that have no technically or economically viable alternatives (e.g., certain medical and research applications, and dental amalgam) (ECCC, 2014). In the case of lamps, rather than introducing a prohibition, the Regulations limit the amount of mercury contained in fluorescent and other types of lamps. The objective of the Regulations is to reduce releases of mercury from products used in Canada to the lowest level that is technically and economically feasible. In February 2018, Canada published a consultation document on proposed amendments to the *Products Containing Mercury Regulations* that aim to fully align Canada with requirements respecting products under the Minamata Convention on Mercury.

3.2.2 Pollution Prevention Actions

The *Notice Regarding Pollution Prevention Planning in Respect of Mercury Releases from Dental Amalgam Waste* was published in 2010 by ECCC. This notice targets dental facilities that have not implemented all of the best management practices set out in the Notice and outlines the requirements to prepare and implement pollution prevention plans for mercury releases from dental amalgam waste.

The *Notice Regarding Pollution Prevention Planning with respect to Mercury Releases from Mercury Switches in End-of-life Vehicles Processed by Steel Mills* was published in 2007 as a Final Notice under Part 4 of CEPA. The Notice targets vehicle manufacturers and steel mills, requiring the preparation and implementation of a pollution prevention plan for the management of mercury switches from end-of-life vehicles, to reduce the release of mercury to the environment (ECCC, 2007).

The *National Strategy for Safe and Environmentally Sound Disposal of Lamps Containing Mercury Act* was enacted in June 2017 and requires the Minister of the Environment to develop a National Strategy for Safe and Environmentally Sound Disposal of Lamps Containing Mercury. Development of the National Strategy is underway in collaboration with provinces, territories, other governments, and other stakeholders, and will be completed by June 2019. The National Strategy may include: the identification of practices for the safe and environmentally sound disposal of those lamps; the establishment of guidelines for facilities where activities involved in the safe and environmentally sound disposal of those lamps are carried out; and, the development of a plan to promote public awareness of the importance of those lamps being disposed of safely and in an environmentally sound manner.

3.2.3 Risk Management Actions

The *Risk Management Strategy for Mercury* provides a comprehensive and consolidated description of the Government of Canada's progress to date in managing the risks associated with mercury. It identifies the objectives, priorities, existing and anticipated actions targeting various industrial sectors, and monitoring programs in place to address the ongoing risks associated with mercury (ECCC and Health Canada, 2010). The ultimate environmental and human health objective for mercury in Canada is

minimizing, and where feasible, eliminating anthropogenic emissions and releases to the environment and minimizing mercury exposure to Canadians. The existing risk management objective for mercury is to minimize releases of mercury from anthropogenic sources into the Canadian environment that is technically and economically feasible.

The *Risk Management Strategy for Mercury-containing Products*, published in 2006, provided a framework for the development of controls to manage mercury-containing products and the environmental effects of mercury used in products (ECCC, 2006). Several strategies were proposed and many were later integrated into the *Products Containing Mercury Regulations*, such as:

- Prohibition of mercury-containing products for which mercury-free alternatives exist
- Prohibition of mercury use in new products, not currently available on the Canadian market, with possible exemptions
- Mercury content limits in products for which mercury-free alternatives do not exist (novelty items excluded)
- Labeling requirements for mercury-containing products.

Remediation of mercury contaminated sediment was undertaken at the [Peninsula Harbour Area of Concern](#) in 2012 and the [St. Clair River Area of Concern](#) from 2002 to 2004.

3.2.4 Monitoring, Surveillance, and Other Research Efforts

Monitoring and measurement for mercury is conducted in water, sediment, air, precipitation, wildlife (including fish and herring gull eggs), and human biomonitoring at sites across Canada, including in the Great Lakes basin. Measurements are collected by ECCC through several different initiatives, including the national Chemicals Management Plan (Government of Canada, 2016). An in-depth assessment of surface waters, surface sediments, and sediment cores is made on a rotational for each of the Canadian Great Lakes annually. Additional water and sediment samples may also be collected from the connecting channels of the annually assessed lake.

Additional monitoring is conducted under regional Great Lakes-specific monitoring and surveillance programs for air, precipitation, herring gull eggs, fish, sediment, and water. The following list highlights select programs being done to acquire mercury data within Canada:

- Human Biomonitoring. In Canada, there are currently no routine Great Lakes-specific human biomonitoring programs to monitor human exposure to persistent chemicals. Therefore, nationwide studies and results of individual epidemiological studies undertaken in the Great Lakes are used to evaluate mercury concentrations in humans in the Great Lakes basin. The Canadian Health Measures Survey (CHMS), initiated in 2007, is a national survey led by Statistics Canada, in partnership with Health Canada and the Public Health Agency of Canada and includes assessment of blood, urine, and hair collected from survey participants for a wide variety of environmental chemicals (Statistics Canada, 2016). The CHMS has recently examined the mean blood concentrations of methylmercury in Canadians and the First Nations Biomonitoring Initiative (FNBI) examined the concentration of inorganic mercury in the blood of Canadian First Nations populations in 2011 (AFN, 2013). Samples were assessed for total mercury in the blood and inorganic mercury in urine. The coefficient of variation for the blood samples was too high to allow for comparison. No comparison could be made between the urine sample sets as more than 40% of the samples were below the limit of detection. However, findings in the urine samples did lead the authors to believe that there may be a greater exposure to inorganic

mercury in the FNBI population than the CHMS population, but further investigation is required (AFN, 2013; Statistics Canada, 2016).

- The Fish Contaminants Monitoring and Surveillance Program (FCMSP) began in 1977 with a focus on monitoring activities in the Canadian waters of the Great Lakes as part of GLWQA. Canada's program activities have been coordinated with those of the United States to ensure basin-wide coverage and to minimize duplication of efforts. The program targets predatory and long-lived fish species that bioaccumulate contaminants. Forage fish, benthos, and plankton are also sampled. Mercury is one of the contaminants of interest analyzed in tissues for the FCMSP.
- The Ontario Ministry of the Environment and Climate Change has monitored mercury in a variety of sport and forage fish in Ontario, Canada since the 1970s in partnership with Ontario Ministry of Natural Resources Forestry (OMNRF) and various other agencies/ institutes. Results are used to produce the [Guide to Eating Ontario Fish](#) well as to analyze trends.
- Canadian Atmospheric Mercury Measurements. Since 1994, considerable atmospheric mercury monitoring and research has taken place across Canada through both ongoing networks and independent research programs. Most monitoring began as independent research programs to measure total gaseous mercury (TGM) in the early 1990s. Realizing the benefits of a community, researchers joined forces to create the Canadian Atmospheric Mercury Measurement Network (CAMNet) in 1994. CAMNet was operated by ECCC from 1994 to 2007, with between 7 and 15 sites across Canada. Later, some of these sites were transferred to the Canadian Atmospheric and Precipitation Monitoring Network (CAPMoN), which still operates these sites today, and to other networks. The remainder of the currently operated ECCC sites are either part of the Northern Contaminants Program (NCP) or are run as part of ECCC measurement programs. As of 2017, these individual programs have been consolidated and fall under Environment and Climate Change Canada – Atmospheric Mercury Monitoring or ECCC-AMM. Currently, there are 12 sites in Canada that collect continuous TGM. As of January 2017, The ECCC-AMM monitors TGM at 12 sites, total particulate mercury (TPM) (termed speciated atmospheric mercury) at 6 sites and wet deposition at 5 sites. There is only one long-term Great Lakes mercury monitoring station in Canada. It is located at the Egbert site, north of Toronto, Ontario.
- The Canadian Mercury Science Assessment (CMSA) was the first comprehensive scientific evaluation and synthesis of mercury in the Canadian environment. The Clean Air Regulatory Agenda (CARA) Mercury Science Program was developed in 2007 to further understand the status of mercury in the Canadian environment and the impact of mercury on Canadian ecosystems and population, and to establish the scientific knowledge base to support regulatory decision-making for mercury. The CMSA was a key outcome of that program. The final document included contributions from more than 230 researchers. This scientific assessment of mercury in Canada has led to the recognition of several detailed priority gaps of knowledge, and has provided scientific recommendations for focus areas and future research, including the integration of mercury impacts on human health, wildlife, and the ecosystem (ECCC, 2016a).

Mercury monitoring and release data results from various programs are centralized in the [Canadian Federal Government Open Data Portal](#). This portal includes data held by the National Atmospheric Chemistry Database (NAtChem) and the National Pollutant Release Inventory (NPRI).

- [NAtChem](#) is a data archival and analysis facility operated by the Science and Technology Branch of ECCC. The purpose of the NAtChem database is to enhance atmospheric research through the archival and analysis of North American air and precipitation chemistry data, including results of research investigations into the chemical nature of the atmosphere, atmospheric processes, spatial and temporal patterns, source-receptor relationships, and long-range transport of air pollutants. The NAtChem Database contains air and precipitation chemistry data from many major regional-scale networks in North America, as well as several major short-term special studies. Contributing networks to NAtChem generally must operate for a period of at least two years, have wide area coverage, and include regionally representative sites (rural and background). The NAtChem Database consists of several smaller databases, one of which contains mercury data. The mercury database contains three types of atmospheric mercury measurements: mercury in precipitation, total gaseous mercury, and mercury speciation in aerosols and gases (ECCC, 2017a).
- [NPRI](#), Canada's PRTR, has tracked releases, disposals, and transfers of mercury (and its compounds) from industrial sources since the NPRI program's inception in 1993. From 1993 to 1999, facilities were required to report mercury once they met the standard NPRI reporting threshold (this threshold was originally based on the thresholds used by the United States TRI). More specifically, during this period facilities were required to report their annual releases, disposals, and transfers of mercury to the NPRI if they met the following reporting criteria:
 - The facility had the equivalent of 10 or more full-time employees; and
 - They manufactured, processed, or otherwise used (MPO) at least 10 tonnes of mercury and its compounds at a concentration of at least 1% by weight.

Starting in 2000, the threshold for the reporting of mercury to the NPRI was lowered to 5 kg MPO at any concentration. According to the data reported to the NPRI for 2015, Canadian facilities in the vicinity of the Great Lakes released a total of 605 kg of mercury (and its compounds); the majority of these releases were reported as releases to the atmosphere. In addition, 34,678 kg of mercury was reported as disposals and 6450 kg were transferred off-site for recycling within this region. In the area surrounding the Great Lakes, the largest mercury releases were reported by facilities in the following sectors: iron and steel mills and ferro-alloy manufacturing, cement manufacturing, sewage treatment, gypsum product manufacturing, non-ferrous metal (except aluminum) smelting and refining, and fossil-fuel electric power generation (ECCC, 2016c).

3.2.5 Canadian Environmental Quality Guidelines and Standards

Canadian Environmental Quality Guidelines (CEQGs) for Mercury have been developed by the Canadian Council of Ministers of the Environment (CCME), which consists of federal, provincial, and territorial environment ministers. CEQGS are nationally endorsed, science-based goals for the quality of atmospheric, aquatic, and terrestrial ecosystems. CEQGS are benchmarks that are recommended as levels that should result in negligible risk to biota, their functions, or any interactions that are integral to sustaining the health of ecosystems. **Table 7** presents CEQGs for mercury in various media.

The Canada-Wide Standards (CWS) Program is a framework under which federal, provincial, and territorial environment ministers address key environmental protection and health risk reduction issues that require common environmental standards across the country. CWS are developed to achieve nationally unified environmental objectives while allowing participating jurisdictions to implement

complementary plans in a way that suits their individual circumstances. The CWS are developed by the CCME. CWS are based on science, but also take into consideration technical feasibility and socio-economic factors. Generally, the CWS consist of a numeric limit and a timeline that sets a deadline for achieving the limit. Implementation plans, progress monitoring, and public reporting are also important aspects of the CWS process. With respect to mercury, CWS have been developed for mercury emissions from base-metal smelters, waste incinerators, coal-fired power plants, mercury-containing lamps, and dental amalgam waste. Implementation plans of each jurisdiction can be found on the [CCME website](#). Brief descriptions of the CWS for mercury emissions are below:

- **Base Metal Smelting.** For base metal smelting facilities, a two-part standard was established to address both existing and new/expanding operations (CCME, 2000). New and expanding facilities should be equipped to meet a guideline of 0.2 g mercury per tonne of finished zinc, nickel, or lead, or 1 g mercury per tonne of finished copper. As of 2009, all but one base metal smelting facilities met the CWS emissions goal of 2 g mercury/tonne of finished metals (CCME, 2010). The single facility that did not meet this standard was not located within the Great Lakes basin and was closed in 2010.
- **Waste Incineration.** Under the CWS, limits for the concentration of mercury in exhaust gas were established for various types of waste incinerators. Larger (processing greater than 120 tonnes of waste per year) facilities must conduct annual stack tests to confirm that the target concentration is achieved. Smaller facilities must make determined efforts, such as an ongoing review of waste diversion options and/or emission control upgrades, to reduce mercury emissions (CCME, 2000). The CWS for hazardous waste incinerators was to achieve a maximum concentration of exhaust gases of 50 µg/Rm³ by 2003. As of 2007, all six hazardous waste incinerators in Canada were in compliance, including the five facilities in Ontario (CCME, 2007).
- **Coal-Fired Electric Power.** CWS consists of two sets of targets: (1) provincial caps on mercury emissions from existing coal-fired electric power generating plants, with the 2010 provincial caps representing a 60% national capture of mercury from coal burned, or 70% including recognition for early action; and (2) capture rates or emission limits for new plants, based on best available control technology, effective as of 2006 (CCME, 2000). In 2014, the capture rate of mercury from coal-generated power plants was 67%, exceeding the CWS goal (CCME, 2016). From 2003 to 2014, total mercury emissions from coal-fired electric plants decreased by more than 75% (CCME, 2016). In January 2017, the Government of Canada released the Pan-Canadian Framework on Clean Growth and Climate Change, and indicated its intention to phase out coal-fired electricity generation to cleaner sources by 2030. To deliver on this intention, the Government of Canada's *Reduction of Carbon Dioxide Emissions from Coal-fired Generation of Electricity Regulations* (2012) and its announced amendments (2016), are expected to generate co-benefits of reducing releases of metals, including mercury, to varying degrees depending on the actions taken by the facilities (e.g., closure, installation of carbon capture and storage technologies, or transition to low-emission fuels). In Ontario, coal-fired electricity production ceased in 2014 resulting in zero release of mercury in the province and into the Great Lakes from locally situated coal-fired plants (CCME, 2016).
- **Mercury-containing Lamps.** The CWS for Mercury-containing Lamps takes a pollution prevention approach by calling for a reduction in the average mercury content of lamps sold in Canada. From a 1990 baseline, the numeric target is a 70% reduction by 2005 and a total reduction of 80% by 2010 (CCME, 2001). By 2004, the average mercury content per lamp had reduced by 73.5% from the 1990 baseline, exceeding the 2005 CWS target (CCME, 2007). Recycling projects

funded by the Ontario Ministry of the Environment collected and recycled more than 14,000 lamps in 2007, capturing more than 400 grams of mercury from within Ontario (CCME, 2007).

- *Dental Amalgam Waste.* The goal of the CWS for Dental Amalgam Waste is to reduce national mercury releases from dental amalgam waste by 95% by 2005 from a 2000 baseline. This CWS promotes the use of best management practices, such as the installation of ISO-certified amalgam traps to achieve the target. It led to the signing of a Memorandum of Understanding by ECCC and the Canadian Dental Association (Canadian Minister of the Environment and Canadian Dental Association, 2002), which focuses on reducing mercury releases from dental facilities. By 2007, 70% of dentists across Canada were using ISO separators, allowing for a 57% reduction in mercury reaching dental wastewaters (CCME, 2007). By 2008, 100% of Ontario dentists had installed amalgam separators to capture waste mercury (US EPA and Canada, 2009). Since by 2007 the 95% target set out in the CWS was not met, Environment and Climate Change Canada took action to achieve the target using pollution prevention planning, through the publication in 2010 of the notice regarding *Pollution Prevention Planning in Respect of Mercury Releases from Dental Amalgam Waste*.

3.3 Binational Actions

3.3.1 Great Lakes Binational Toxics Strategy

The [Great Lakes Binational Toxics Strategy](#) (GLBTS) was a binational P2 endeavor from 1997 to 2007 that focused on the virtual elimination of mercury and mercury-containing compounds, among other persistent, bioaccumulative, and toxic chemicals. The Strategy outlined a four-step process for addressing recognized knowledge gaps regarding the Great Lakes region: (1) gather information; (2) analyze current regulations, initiatives, and programs that manage or control substances; (3) identify cost-effective options to achieve further reductions; and (4) implement actions to work toward the goal of virtual elimination (US EPA and ECCC, 1997).

3.3.2 Lakewide Action and Management Plans

Additional binational mercury pollution prevention actions have been initiated through LAMP programs for each Great Lake. LAMPs are plans of action to assess, restore, protect, and monitor the ecosystem health of each Great Lake (US EPA, 2004; US EPA, 2016d). As an example, the Lake Superior Zero Discharge Demonstration Program (ZDDP) was established in 1991 by the Lake Superior Binational Program (LSBP) with a goal of eliminating nine persistent, bioaccumulative, and toxic pollutants, including mercury, from Lake Superior by 2020 (Lake Superior Partnership, 2016). Under the Lake Superior ZDDP, an inventory of mercury (equipment and waste containing mercury) in the Lake Superior basin has been updated approximately every 5 years since 2000. Other LAMPs are in the process of reviewing their monitoring priorities, which may include CMCs in the future.

3.3.3 Great Lakes Regional Collaboration

The Great Lakes Regional Collaboration (GLRC) was a cooperative effort between the Governors and Premiers of the Great Lakes States and Provinces. Members of the GLRC include the Great Lakes Interagency Task Force, the Council of Great Lakes Governors, the Great Lakes and St. Lawrence Cities Initiative, the Great Lakes Native American Tribes, and the Great Lakes Congressional Task Force. The GLRC has produced two key strategies for Mercury:

- *Great Lakes Mercury in Products Phase Down Strategy (2008).* Great Lakes Mercury in Products Phase Down Strategy was developed in response to the GLRC Strategy to Restore and Protect

the Great Lakes. The phase down strategy established timelines for full phase-outs of mercury-added products by 2015.

- *Great Lakes Mercury Emission Reduction Strategy (2010)*. The Great Lakes Mercury Emission Reduction Strategy was developed in response to the GLRC Toxic Pollutants Initiative, which called for the development of a basin-wide mercury emission strategy designed to phase out the use of mercury and provide for mercury waste management. The Strategy was not intended to be a comprehensive summary of actions to reduce mercury releases to the Great Lakes; rather, it addressed mercury air emissions with a primary goal to reduce mercury emissions from new and existing sources whose mercury emissions have not been regulated, and from sources where regulations have been implemented but additional cost-effective reductions can be achieved. The Strategy was finalized in December 2010 and presented 34 recommendations for reducing mercury emissions in seven source sectors, as well as cross-cutting strategies and actions to track progress of implementation (GLRC, 2010). By 2014, approximately 75% of the recommendations had been completed or were continuing across the eight Great Lakes States (GLRC, 2014).

3.3.4 National Atmospheric Deposition Program/Mercury Deposition Network

The Mercury Deposition Network (MDN) is a long-term mercury wet-deposition monitoring network of the National Atmospheric Deposition Program (NADP). NADP is a cooperative program between federal, state, provincial, academic institutions, indigenous peoples, and private organizations. MDN provides a long-term record of total mercury concentration and deposition in precipitation across the United States and Canada. The MDN began measuring total mercury in precipitation in 1996. It now includes more than 100 sites. All MDN sites follow standard procedures and have uniform precipitation chemistry collectors and gages to allow for direct comparison between sampling sites. While total mercury in precipitation is measured at all locations, methylmercury is monitored at select sites (NADP, 2011).

3.3.5 Cooperative Science and Monitoring Initiative

One aspect of the GLWQA is the establishment of a Cooperative Science and Monitoring Initiative (CSMI) Task Team through Annex 10. The charge of the CSMI is to implement a joint United States-Canadian effort to provide environmental and fishery managers with the science and monitoring information necessary to make management decisions for each Great Lake. A five-year rotating cycle in which the lakes are visited one per year is followed by an intensive CSMI field year. By studying one Great Lake per year, science and monitoring activities can focus on information needs not addressed through routine agency programs, and specific science assessments can be coordinated. Individual Lake Wide Partnerships identify science needs according to the CSMI schedule, and the Task Team implements these recommendations, when appropriate.

3.3.6 State of the Great Lakes Reporting

Between 1994 and 2011, the US EPA and ECCC coordinated reporting efforts under the State of the Lakes Ecosystem Conferences (SOLEC) (ECCC and US EPA, 2011). Since then, the SOLEC conference has been replaced with scientific confirmation webinars to review a triennial assessment and report from the governments of Canada and the United States (ECCC and US EPA, 2017). These reports provide decision-makers and scientists the opportunity to receive comprehensive and current information on the state of the Great Lakes through three main elements:

- Interactive scientific webinars convened triennially for Great Lakes stakeholders to review assessments and reports and provide additional information or interpretation.

- State of the Lakes reports incorporating a suite of indicators of the status and trends of Great Lakes ecosystem components developed by Great Lakes agencies, organizations, and other stakeholders as the basis for the triennial assessments.
- A comprehensive State of the Great Lakes technical report based on findings from the indicators.

3.4 International

Both the United States and Canada participate in a number of initiatives and partnerships established at the international level to limit the availability, use, release, and overall number of mercury sources.

3.4.1 United Nations Economic Commission for Europe, Convention on Long-Range Transboundary Air Pollution

The 1998 Aarhus Protocol on Heavy Metals to the Convention on Long-range Transboundary Air Pollution (LRTAP) aims to reduce emissions from industrial sources and products for three harmful metals: mercury, cadmium, and lead. Stringent limits were established for emissions from stationary sources and best available techniques were suggested for use. The protocol proposed management measures for mercury in batteries, electrical components, measuring devices, fluorescent lamps, dental amalgam, pesticides, and paint (ECE, 2014; UNECE, 2017). In 2012, the protocol was amended to encourage ratification from UNECE countries with economies in transition.

3.4.2 United Nations Environment Program, Minamata Convention on Mercury

The Minamata Convention on Mercury (Minamata Convention) currently has 128 signatory countries and as of February 1, 2018 has been ratified by 88 Parties (<http://mercuryconvention.org/>). The United States and Canada are both Parties, with the United States having joined in 2013 and Canada in 2017. The Convention is the first new global convention related to environment and health in close to a decade. The Minamata Convention builds upon the 1998 Protocol on Heavy Metals to raise the profile of mercury to a global level (UNECE, 2017). The Convention covers the entire lifecycle of mercury pollution resulting from human activities and includes banning new mercury mines, phasing-out existing ones, prohibiting the majority of mercury-containing products and mercury processes, reducing artisanal and small-scale gold mining, and reducing emissions and releases.

The Convention entered into force in August 16, 2017, and the first meeting of the Conference of the Parties (COP1) to the Minamata Convention took place in September 2017 in Geneva, Switzerland. Parties to the Convention will have committed to the implementation of specific measures to control mercury pollution:

- Control, and where feasible reduce mercury air emissions from coal-fired power plants, coal-fired industrial boilers, certain non-ferrous metals smelting and roasting operations (lead, zinc, copper and industrial gold), waste incineration, and cement clinker production, including use of best available techniques and best environmental practices for new and substantially modified sources.
- Phase out mercury in the majority of products (e.g., batteries, switches, lights, cosmetics, pesticides, and measuring devices) and create initiatives to reduce the use of mercury in dental amalgam.
- Phase out or reduce the use of mercury in manufacturing processes such as chlor-alkali production, vinyl chloride monomer production, and acetaldehyde production.

- Address the supply and trade of mercury, its interim storage and final disposal, and develop strategies for contaminated sites.

The Minamata Convention will enable information and assistance exchange, public awareness, research, and monitoring. Parties of the Convention will be required to report on measures taken to implement certain provisions. The Convention will be periodically evaluated to assess its effectiveness at meeting its objective of protecting human health and the environment from mercury pollution (UN Environment, 2017a).

3.4.3 UNEP Global Mercury Partnership

In February 2005, the UNEP Governing Council established the Global Mercury Partnership, a voluntary multi-stakeholder partnership aimed at reducing use and emissions of mercury globally to protect human health and the environment from the releases of mercury and its compounds. The United States was a catalyst in the formation of the Partnership. The US EPA serves as lead for the Mercury Cell-Chlor-Alkali and Product Partnership Areas, and co-chairs the Partnership Advisory Group, which provides advice and guidance to the Global Mercury Partnership to encourage the work of the partnership areas.

The Partnership is divided into seven areas with the following leads or co-leads:

- Mercury Emissions from Coal – International Energy Agency Clean Coal Center
- Mercury Air Transport and Fate Research – Consiglio Nazionale delle Ricerche (CNR) Institute for Atmospheric Pollution, Italy and Biodiversity Research Institute (BRI)
- Mercury Waste Management – Government of Japan
- Mercury Cell Chlor-Alkali Production – Government of the United States (US EPA), United Nations Industrial Development Organization (UNIDO)
- Artisanal and Small-Scale Gold Mining – Natural Resources Defense Council (NRDC), United Nations Industrial Development Organization (UNIDO)
- Mercury Supply and Storage – Government of Spain, Government of Uruguay
- Mercury in Products – Government of the United States (US EPA).

Canada is a member of the Mercury Air Transport and Fate Research as well as the Mercury Emissions from Coal Partnerships.

3.4.4 Arctic Contaminants Action Plan and the Arctic Monitoring and Assessment Program under the Arctic Council

Arctic Contaminants Action Plan (ACAP) and Arctic Monitoring and Assessment Program (AMAP) are bodies of the Arctic Council, an intergovernmental forum whose members include Canada, Russia, Norway, Denmark, Iceland, the United States, Sweden, and Finland. US EPA co-chairs the Mercury Project Steering Group under ACAP. In 2011, this group published a list of [nine recommendations](#) for reducing mercury in the artic (AMAP, 2011).

3.4.5 The Commission for Environmental Cooperation

The Commission for Environmental Cooperation (CEC) is a cooperative agreement between the three countries of North America: Canada, Mexico, and the United States. The CEC's mission is to facilitate collaboration and public participation in conservation, protection, and enhancement of the North

American environment, specifically in the context of increasing economic, trade, and social links among the three countries. Mercury is considered persistent, bioaccumulative, and toxic by the CEC. Thus, the CEC Council has prepared a North American Regional Action Plan (NARAP) to determine the level of contamination in the environment and in humans, to follow the trends over time, and to support mercury monitoring needs (CEC, 2015).

4 Gap Analysis

4.1 Gaps and Needs for Action

The largest contributions of mercury loadings to the Great Lakes basin are through electricity generation facilities within the United States, and from long-range transport. In each case, air monitoring is critical for assessing trend analysis, source attribution and predictive modeling. Mercury in Great Lakes air, precipitation, water, sediment, fish, and wildlife has been monitored to some degree since the 1960's; however, uniform long-term monitoring is lacking in areas of the Great Lakes basin. Data collected by federal, state, provincial, Tribal/First Nations/Métis, and other governmental agencies or programs need to be consistent and of uniform quality.

There is an overall lack of knowledge in how the changing climate may impact the mercury cycle. Climate can impact physical characteristics and functions of the ecosystem; this affects all processes in the biogeochemical cycle of mercury (ECCC, 2016b). Of concern is the rate of methylmercury formation within the environment, and its corresponding ecological risk. Additional topics that have been previously identified as mercury cycle knowledge gaps include (ECCC, 2016b):

- Limited information on the processes driving emission of mercury from various surfaces and water bodies;
- Lack of chemical identification of different species of atmospheric mercury and quantification of their deposition to surfaces;
- Lack of knowledge of the impact of acidity, temperature, and organic matter collectively on methylmercury production and bioaccumulation in freshwater aquatic systems;
- Insufficient information on methylmercury levels and production in the freshwater environment;
- Insufficient knowledge of factors promoting methylmercury in Great Lakes food webs and potential impacts of invasive species; and
- Lack of information on the fate and transport of mercury within terrestrial ecosystems.

There is also a gap in the ability of models to accurately predict the relationship between mercury emissions and fish concentrations due to the intrinsic complexity of atmospheric mercury chemistry and mercury methylation pathways and processes. Additional emissions data and monitoring data from both biotic and abiotic systems are needed to enhance the utility of models to effectively predict fish populations within specific areas that are at risk due to mercury exposure (ECCC, 2016b). Such models would be useful to provide a comprehensive evaluation of how well existing regulatory programs are working to limit and reduce mercury impacts to the Great Lakes, another current gap. Further development of innovative tools like isotopic mercury ratios will help with a weight-of-evidence approach to evaluation of sources to the Great Lakes.

Within the United States, monitoring to assess the effects of the Mercury Export Ban Act of 2008 (MEBA) is needed. MEBA is intended to reduce the availability of elemental (metallic) mercury in local and global markets. MEBA amends TSCA to prohibit the export of elemental mercury from the United States effective January 1, 2013. By reducing the supply of elemental mercury in commerce, MEBA seeks to reduce the use of mercury in artisanal mining and other commercial purposes globally. However, there is a concern that higher prices for recycled mercury will result in primary mercury mining (Bender and Narvaez, 2016). Primary mercury mining produces very significant releases of mercury to the environment, and is an overall less environmentally sound method of meeting global mercury demand. Therefore, careful evaluation and consideration should be given to the global impact of export bans.

There have been a number of studies examining the relative health benefits of fish consumption compared with the risks of mercury exposure (Ginsberg and Toal, 2009; US FDA, 2014; Ginsberg, 2016; Raymond et al., 2017). There is, however, an ongoing need to assess the balance between the nutritional benefits of fish consumption and the risk of methylmercury exposure as additional information becomes available particularly since the relationship between methylmercury exposure and other diseases/conditions is not fully understood or appreciated.

4.2 Exceedances of or Non-compliance with Environmental Quality Guidelines

While many environmental quality guidelines exist for mercury in Canada and the United States, not all sample types are regularly recorded. From the available information, the offshore waters in the Great Lakes are below the water CEQG for the protection of aquatic life; however, sediment concentrations within some lakes (Lakes Erie and Ontario) regularly exceeded the sediment CEQGs (**Table 7**) (ITT, 2015).

The concentration of total mercury in whole Lake Trout from the Great Lakes has been shown to be below levels of concern for the health of fish consuming wildlife that were established in the 1987 GLWQA. However, mercury levels in fish are still sufficient to trigger fish consumption advisories in all five of the Great Lakes (**Figure 8**). Long-term mercury trends in walleye and largemouth bass fillets from the Great Lakes indicate decreasing concentrations, however, fish fillet concentrations still exceed US EPA mercury criteria for the protection of human health ($0.3 \mu\text{g/g}$ wet weight) and Canadian restrictions for sport fish consumption by sensitive populations ($0.026 \mu\text{g/g}$ wet weight) (Evers et al., 2011; Monson et al., 2011; OMOECC, 2013; ITT, 2015).

5 Risk Mitigation and Management Options to Address Gaps

The options highlighted herein represent both new and the continuation of current risk mitigation and management actions to address the identified gaps. Implementation may result in measurable (either qualitatively or quantitatively) human health and/or environmental benefits, or enhanced understanding of mercury sources, fate, and human health/environmental effects.

5.1 Regulations, and Other Risk Mitigation and Management

In accordance with the recently enacted Lautenberg Act amendments to TSCA, the US EPA is required to establish a mercury inventory for supply, use, and trade in the United States, and requires reporting by any person who manufactures mercury or mercury-added products or otherwise intentionally uses mercury in a manufacturing process. This information could be used to periodically update the mercury inventory, identify manufacturing processes or products that use mercury, and recommend actions to reduce mercury use.

The Lautenberg Act also augmented the ban on exports of elemental mercury under MEBA to include certain listed mercury compounds under TSCA. However, the effectiveness of such regulatory programs

has yet to be comprehensively evaluated from a global perspective. Such evaluation of existing regulatory programs could lead to increased efficiencies and an overall decrease in mercury emissions.

In 2017, Canada added mercury to the Export Control List and amended the associated Export of Substances on the Export Control List Regulations under CEPA. These amendments require exporters of mercury and products containing mercury to notify the Minister prior to export and set out the limited purposes for which export may occur (ECCC 2018). Comprehensive restrictions apply to exports of elemental mercury at concentrations of 95% or more by weight. Until recently, mercury exports were not controlled or prohibited, except as hazardous waste or hazardous recyclable material regulated under the *Export and Import of Hazardous Waste and Hazardous Recyclable Material Regulations*.

In 2010, the Great Lakes Regional Collaboration (GLRC), a binational task force, produced the Great Lakes Mercury Emission Reduction Strategy (GLRC, 2010). The strategy includes more than 34 recommended regulatory and voluntary actions to further control mercury pollution, including the following:

- *Require best available control technology for new and modified sources.* The GLRC strategy recommended that all states require Best Available Control Technology for new and modified sources if they annually emit 10 pounds of mercury (or less, at the state's discretion).

Summary of Regulations and Other Risk Management Strategy Options

- Evaluate the effectiveness of existing regulatory programs to ensure maximum efficiency and overall positive implications on a global scale (Canada)
- Evaluate the effectiveness of existing emissions regulatory programs for addressing mercury pollution (US)
- Review and update actions to match current scientific understanding and regional context (Canada and US)
- Identify manufacturing processes or products that intentionally add mercury (US)
- Continue to reduce mercury emissions resulting from coal-fired generation of electricity (Canada)
- Continue implementation of domestic regulations and other risk management activities for mercury (Canada and US)
- Develop the National Strategy for Safe and Environmentally Sound Disposal of Lamps Containing Mercury (Canada)
- Continue remediation of mercury-contaminated sites and sediments (Canada and US)
- Amend the Products Containing Mercury Regulations to further reduce mercury in products (Canada)

5.2 Compliance Promotion and Enforcement

While there are many regulations in effect for mercury and mercury emissions, these regulations need to be effectively communicated to the appropriate entities for full compliance. These regulations were discussed in Section 3.

Internationally, both the United States and Canada are Parties to the Minamata Convention. Canada and the United States should continue to support international initiatives to reduce mercury emissions and prevent release of mercury into the environment. Implementation of the Minamata Convention is essential and guidance adopted at COP1 will help address the next phase of global implementation.

Summary of Compliance Promotion and Enforcement Strategy Options

- Promote compliance with domestic and international mercury activities and initiatives (Canada and US)
- Continue implementation of respective obligations of the Minamata Convention on mercury (Canada and US)

5.3 Pollution Prevention

User-friendly documents are needed to educate and engage the general public in efforts to reduce the potential for mercury release or exposure. User-friendly materials targeting specific public audiences may be appropriate to aid in preventing mercury pollution from being incorporated into general solid waste streams, and providing awareness regarding potential mercury sources of health hazards. Outreach/education is also needed to ensure affected populations have an awareness and understanding of existing fish consumption advisories. Key elements of outreach and education include knowledge of where to obtain the most timely and relevant information and how to apply the advisories to individuals (e.g., young children, pregnant women).

US EPA's TRI database can be used to track progress in reducing waste generation. The TRI database should be maintained and leveraged to maximize P2 activities being conducted by industries in the Great Lakes region. Highlighting pollution prevention successes in the Great Lakes basin may be beneficial in increasing awareness, coordinating P2 efforts in similar sectors throughout the basin, and furthering the reduction of mercury in the environment. Waste reduction success stories may be noted in region-specific journals, websites, and/or at conferences.

Canada's NPRI database tracks mercury emissions. Industries can see their emissions and releases decreasing over time as they implement P2 activities.

Summary of Pollution Prevention Strategy Options

- Enhance public outreach and educate the public and facility staff on potential sources of mercury and proper actions to follow when handling mercury containing products (Canada and US)
- Enhance public outreach and educate the public on how to obtain and implement site-specific fish consumption advisories (Canada and US)
- Encourage industries to track their P2 activities and efforts in the National Pollutant Release Inventory (NPRI) or Toxics Release Inventory (TRI), or via P2 promotion activities (fact sheets, case studies) (Canada and US)
- Highlight pollution prevention successes (Canada and US)
- Implement best available techniques and best environmental practices for new and substantially modified sources (Canada and US)

5.4 Monitoring, Surveillance, and Other Research Efforts

Numerous monitoring, surveillance, and remediation activities have been conducted since the 1970s. However, due to the complexity of mercury and methylmercury interactions in the environment, a clear understanding of the long-term interactions is lacking. Domestically, both Canada and the United States should continue to assess and, where appropriate, remediate mercury-contaminated sites and sediments related to historical sources and evaluate the effectiveness of remediation activities.

Additional research is needed to fill knowledge gaps with respect to methylation dynamics and the differential impacts of mercury in nearshore versus offshore environments. Monitoring data, particularly for air, is needed for inclusion in models to track long-range atmospheric transport, the rate of methylmercury formation in the environment, and the implications of climate differences for the overall mercury cycle. These data will also help assess the effectiveness of the implemented regulations. The rate of methylmercury formation corresponds to ecological risks and risk associated with human consumption of Great Lakes fish populations. Therefore, a clear understanding is imperative.

The development of a cost-effective and useful means of collecting mercury concentrations from a variety of sources as part of a comprehensive mercury monitoring network (eg. MercNet) is essential. Implementation of a passive sampler capable of monitoring total mercury levels in air and development of passive samplers for monitoring speciated mercury compounds could be used in conjunction with active sampling to better understand the spatial distribution and behavior of mercury in the Great Lakes and the region as a whole (Huang et al. 2014, McLagan et al. 2016). Source tracking for areas of localized mercury contamination may be a need in the future. Efforts such as Project Trackdown, which uses a multimedia weight-of-evidence approach for tracing PCB sources in the Great Lakes, may be a model system for future mercury studies (Benoit et al., 2016).

Reports on the current state of mercury contamination and the ecological and human health risks that mercury poses have been published in peer-reviewed journals, websites, and social media. Each form of reporting is designed to target specific audiences to maximize the application of the results. Results of future monitoring efforts should continue to be published in multiple formats to effectively communicate changes observed in the Great Lakes region.

While the ultimate success of the strategy depends on the efforts of the Great Lakes community, it is suggested that the strategy and its implementation progress should be reviewed on a regular basis and reported through the Progress Report of the Parties.

Summary of Monitoring, Surveillance, and Other Research Strategy Options

- Continue monitoring mercury in environmental media in the Great Lakes (air, precipitation, sediment, fish, and other wildlife) and publish results in a variety of publications (e.g., online and open data portals, government reports and scientific journals) to maximize the intended audience (Canada and US)
- Continue efforts to update and maintain mercury emissions inventories in a manner such that regional and global emissions can be tabulated (Canada and US)
- Conduct additional research on methylation dynamics and the differential impacts of mercury in nearshore versus offshore environments (US)
- Enhance existing models to track long-range atmospheric transport and the rate of methylmercury formation in the environment and its corresponding ecological risk (Canada and US)
- Develop cost-effective, reliable and effective tools (e.g., passive samplers) for collecting long-term mercury multi-media monitoring data (Canada and US)
- Develop and populate a structured data system to track mercury sources, manifests, waste, and products (Canada and US)

5.5 Domestic Water Quality

Domestic waters include all water used for indoor and outdoor household purposes. There is a need for reviewing existing standards to ensure that they are based on the latest science, to assist states and Ontario (and other Canadian provinces and territories) in identifying areas where standards are exceeded. Compliance with the ambient water quality guidelines of both countries will ensure drinking water is protected since the ambient criteria are well below the drinking water standards.

Summary of Domestic Water Quality Strategy Options

- Review and update existing domestic water quality guidelines and standards, if necessary (Canada and US)

6 Conclusions

Under Annex 3 of the GLWQA, mercury has been identified as a CMC that originates from anthropogenic sources. The documented impacts of mercury exposure of fish and wildlife in the Great Lakes region are significant (Evers et al., 2011). Elevated mercury levels have been detected in Great Lakes biota (e.g., birds, fish, mammals), at all levels of the food web (e.g., plankton, fish, loons), and across many different habitat types (e.g., lakes, wetlands, streams, forests) throughout the region. Much of the point-source mercury pollution has been controlled, allowing partial recovery of the Great Lakes, as shown by lower

mercury concentrations in lake sediments in the lower Great Lakes (e.g., Lake Ontario) and in declines in fish mercury since the years of peak pollution.

There has been significant international, national (United States and Canada), and regional efforts to reduce mercury emissions to the environment since the 1980s. Binational efforts have made significant strides in regulating industrial mercury emissions and in cleaning historic mercury-contaminated sites (US EPA, 2016a). However, mercury is still detected at elevated concentrations in soils, sediment, water, air, biota tissues, and wastes throughout the Great Lakes basin, such that fish consumption advisories due to mercury concentrations are in effect in all five Great Lakes (**Table 7**). Although there are some emitting sectors this is primarily attributable to legacy accumulations from historical emissions and discharges in the U.S. and Canada as well as long-range atmospheric transport.

Under the GLWQA, ECCC and the US EPA have a mutual goal for virtual elimination of mercury emissions originating from human activities in the Great Lakes region. This goal will be achieved through a variety of programs and actions with a primary emphasis on pollution prevention. Binational cooperation is needed to coordinate monitoring and surveillance efforts, maximize research initiatives, and cost-effectively monitor and track mercury concentrations in multiple media (wastes, soil, water, air, tissues, etc.). A broad audience of Great Lakes stakeholders who are committed to protecting and restoring the Great Lakes ecosystem is encouraged to implement the risk mitigation and management options outlined in this document. Continued progress in seeking novel ways and/or improving upon existing ways to mitigate and manage mercury risks will improve the health of the ecosystem and residents of the Great Lakes basin, and will preserve the quality of the Great Lakes for future generations.

7 Figures




Figure 1. The Mercury Cycle. Source: Evers et al. (2011)




Figure 2. Decline in Mercury Emissions 1990-2005. Source: Evers et al. (2011)




Figure 3. Trends in National Emissions Inventory (NEI) Mercury emissions (tons) Source: United States EPA 2014 NEI version 2 Technical Support Document (TSD)




Figure 4. Historical and Projected Canadian Mercury Air Emissions Trends. Source: ECCC (2016b)




Figure 5. Seven-year mean annual mercury wet deposition based on NADP/MDN monitoring data.
Source: Evers et al. (2011)




Figure 6. Spatial Distribution of Mercury in Great Lakes Sediments. Inset is Lake St. Clair Corridor.
Source: State of the Lakes Technical Report (2017)




Figure 7. Long-Term Mercury Trends in Chinook Salmon, Coho Salmon, Lake Trout, Lake Whitefish and Walleye in the Great Lakes. Source: State of the Lakes Technical Report 2017.




Figure 8. Great Lakes Fish Consumption Advisories for Mercury. Source: Evers et al. (2011)

8 Tables

Table 1. Physical and Chemical Properties of Mercury and Select Compounds.

Property	Mercury	Mercuric Chloride	Mercuric Oxide	Mercury Sulfide	Methylmercury Chloride	Dimethyl Mercury
Chemical Symbol	Hg	HgCl ₂	HgO	HgS	CH ₃ HgCl	(CH ₃) ₂ Hg
Physical State	Liquid	Crystalline solid	Crystalline powder or scales	Solid	NA	Liquid
Molecular Weight (g/mol)	200.59	271.49	216.59	232.65	248.69	230.66
Color	Shiny, silvery-white	White	Red, Orange-red, orange-yellow	Red or black	NA	Colorless
Crystal Structure	Rhombohedral	Rombic or Orthorombic	Orthorombic	NA	NA	None
Density, g/cm ³	13.5					
Melting Point, °C	-39	277	Decomposition @ +500	Sublimation 584	Sublimation 167	NA -S
Boiling Point, °C	357 @ 1 atm.	303 @ 1 atm.	NA S	NA S	NA S	96 @ 1 atm.
Vapor Pressure (Pa)	0.180 @ 20°C	8.99 × 10 ⁻³ @ 20°C	9.20 × 10 ⁻¹² @ 25°C	NA S	1.76 × 10 ⁻¹² @ 25°C	8.3 × 10 ³ @ 25°C
Water Solubility (g/L)	49.4 × 10 ⁻⁶ @ 20°C	66 @ 20°C	5.3 × 10 ⁻² @ 25°C	~ 2 × 10 ⁻²⁴ @ 25°C	~ 5 to 6 @ 25°C	2.95 @ 24
Henri's Law Coefficient (Pa m ³ mol ⁻¹)	729 @ 20°C	3.69 × 10 ⁻⁵ @ 20°C	3.76 × 10 ⁻¹¹ @ 25	NA S	1.6 × 10 ⁻⁵ @ 25°C and pH = 5.2	646 @ 25°C
	0.32 @ 25°C					0.31 @ 15°C
	0.18 @ 5°C					0.15 @ 0°C
Octanol-Water Partition Coefficient (dimensionless)	4.2	0.5	NA S	NA S	2.5	180

Sources: Schroeder and Munthe (1998); Kim et al. (2016),

~ - approximately; atm. – atmosphere; NA – not available

Table 2. *Estimated Amount of Elemental Mercury in Mercury-Added Products in the United States (2013)*

PRODUCTS/COMPONENTS	TOTAL MERCURY SOLD IN THE UNITED STATES (METRIC TONS)			
	2001	2004	2007	2013
Switches and Relays	52.44	46.97	27.91	17.7 (2010)
Dental Amalgam	27.91	27.57	14.95	14.5
Thermostats	13.27	12.85	3.50	not reported
Lamps	9.22	8.67	9.64	4.7
Miscellaneous	4.64	2.18	2.52	2.5
Batteries	2.68	2.30	1.88	<0.1
Other products*	5.46	4.35	2.32	0.6
Total (approximate)	115.6	104.51	62.78	40.1

*Chemicals, sphygmomanometers, thermometers, manometers, and barometers

Source: Carpenter et al. (2011 IMERC, US EPA 2017c)

Table 3. *Sources of Global Mercury Supply, 2015.*

Mercury Supply Sector	Range (metric tons)
Dedicated mercury mining	1630–2150
By-product mercury from non-ferrous metal sector	440–775
Recycled/re-use mercury from chlor-alkali facilities	370–450
Recycled mercury—others	1040–1410
Commercially available mercury stocks	As needed (+)
Range of Total	3480–4785+

Source: United Nations Environment Program (2017c)

Table 4. *Total Mercury Emissions in Canada by Sector, 1990-2015.*

Sector	Total Emissions per Year (tons)					
	1990	1995	2000	2005	2010	2015
Non-ferrous Metal Mining and Smelting	24.9	4.7	1.9	1.7	0.5	0.2
Electric Power Generation	2.3	2.0	4.0	2.1	1.6	0.7
Cement and Concrete industry	0.5	0.4	0.4	0.2	0.3	0.3
Iron and Steel Industries	0.9	1.0	1.0	0.7	0.4	0.7
Waste Sectors	3.8	4.3	2.1	1.4	1.3	0.7
Other	2.9	2.4	1.7	1.2	1.2	1.8
Total Emissions	35.3	14.7	11.2	7.3	5.3	4.4
% per year decrease since 1990		58.4	68.3	79.3	85.0	87.5

Source: ECCC (2016b)

Table 5. United States Federal Statutes Authorizing Mercury Regulations.

Regulation	Summary of Elements
Frank R. Lautenberg Chemical Safety for the 21st Century Act on June 22, 2016	Amended TSCA to include additional objectives in the Act specific to mercury and mercury compounds (https://www.epa.gov/assessing-and-managing-chemicals-under-tsca/frank-r-lautenberg-chemical-safety-21st-century-act-5)
Mercury Export Ban Act (MEBA) of 2008	This law focuses on reducing the availability of elemental (metallic) mercury from local and global markets through a reduction in supply. (https://www.congress.gov/110/plaws/publ414/PLAW-110publ414.pdf)
Mercury-Containing and Rechargeable Battery Management Act of 1996	This law phases out the use of mercury in batteries and provides for cost-effective means to dispose of regulated batteries, including used nickel cadmium (Ni-Cd) batteries and small sealed lead-acid (SSLA) batteries. (https://www.epa.gov/sites/production/files/2016-03/documents/p1104.pdf)
Toxic Substances Control Act (TSCA) §6(f) – 15 U.S.C. 2605(f)	Title 1 of TSCA prohibits the sale, distribution, or transfer of elemental mercury by Federal agencies, state or local agencies, or any private individual or entity except for the purposes of storage. (https://www.epa.gov/enforcement/toxic-substances-control-act-tsca-and-federal-facilities)
Clean Air Act (CAA) 42 U.S.C. § 7401 et seq. (1970) Section 112 and Section 129	Section 112 authorizes US EPA to regulate hazardous air pollutant emissions, including mercury, from new and existing stationary sources. Section 129 provides authority to address emissions of pollutants, including mercury, from new and existing solid waste combustion units. (https://www.epa.gov/laws-regulations/summary-clean-air-act)
Clean Water Act (CWA) 33 U.S.C. § 1251 et seq. (1972)	Establishes the basic structure for regulating discharges of pollutants into the waters of the United States and regulating quality standards for surface waters. Authorized the National Pollutant Discharge Elimination System (NPDES), wastewater standards for industry, and water quality standards for contaminants within surface waters. (https://www.epa.gov/laws-regulations/summary-clean-water-act)
Emergency Planning and Community Right-to-Know Act (EPCRA) Section 313	Requires industrial and federal facilities to report mercury emissions through the Toxics Release Inventory (TRI) Program. (https://www.epa.gov/sites/production/files/documents/2001hg.pdf)
Resource Conservation and Recovery Act (RCRA) 40 CFR Parts 262 - 265	Mercury wastes are manifested using the RCRA Uniform Hazardous Waste Manifest. Individual states are largely responsible for implementing the RCRA program and therefore, some states may have stricter requirements. (https://www.epa.gov/laws-regulations/summary-resource-conservation-and-recovery-act)
Safe Drinking Water Act (SDWA) 42 U.S.C. § 300f et seq. (1974)	Authorizes US EPA to establish minimum standards to protect all waters actually or potentially designed for drinking use. Requires all owners or operators of public water systems to comply with primary (health-related) standards. PCB maximum contaminant level goal (MCLG): 2 ppb; maximum contaminant level (MCL): 2 ppb. Individual states have the primary responsibility for enforcing drinking water standards. (https://www.epa.gov/laws-regulations/summary-safe-drinking-water-act)

Table 5. (continued) United States Federal Statutes Authorizing Mercury Regulations.

Regulation	Summary of Elements
Comprehensive Environmental Response, Compensation and Liability Act (CERCLA) or “Superfund” 42 U.S.C. § 9601 et seq. (1980)	Gives the Federal government authority to respond to mercury emergencies and clean up uncontrolled or abandoned hazardous waste sites (US EPA, 2015c). (https://www.epa.gov/laws-regulations/summary-comprehensive-environmental-response-compensation-and-liability-act)
Effluent Limitation Guidelines and Standards for the Dental Category 40 CFR 441	This rule requires dental offices to use amalgam separators and two best management practices recommended by the American Dental Association (ADA). This rule went into effect on July 14, 2017 for all new dental facilities, while pre-existing sources must comply by July 2020 (US EPA, 2017a). (https://www.federalregister.gov/documents/2017/06/14/2017-12338/effluent-limitations-guidelines-and-standards-for-the-dental-category)

Source: US EPA (2017b)

Table 6. Summary of Great Lakes States Mercury Programs from a 2011 Survey.

State	Overall Mercury Action Plan	Inventory Mercury Air Emissions Sources	Mercury Monitoring – Stack Testing	Mercury Monitoring – Air Deposition	Mercury Monitoring – Fish Consumption Advisory	Programs to Manage Mercury – Containing Products	Mercury Dental Programs	State Requirements for Mercury Switch Recovery	Participant in National Vehicle Mercury Switch Recovery Program	Statewide TMDL	Multistate TMDL
Illinois			X	X	X	X	X	X	X		
Indiana	X	X	X	X	X	X	X	X	X		
Michigan	X	X	X	X	X	X	X		X	Draft	
Minnesota	X	X	X		X	X	X		X	X	
New York	X	X	X	X	X	X	X		X		X
Ohio		None	X	X	X	X	X		X		
Pennsylvania*				X	X	X			X	X	
Wisconsin		X		X	X	X	X		X		

Source: Quicksilver Caucus (2012)

* Pennsylvania information was gleaned from the Pennsylvania Department of Environmental Protection (www.dep.pa.gov).

Table 7. United States Standards and Canadian Environmental Quality Guidelines for Mercury and Methylmercury and Average Great Lakes Concentrations.

Medium/Guideline	United States Standards	Canadian Environmental Quality Guideline	Average Great Lakes Concentration	Comment/Source
Mercury				
Drinking Water	2.0 µg /L	1.0 µg/L		US MCL per the Safe Drinking Water Act (ITT, 2015); Health Canada (1986)
Water Quality Guidelines for the Protection of Aquatic Life		26 ng/L	0.24-0.54 ng/L	CCME (1999 and updates); ITT (2015)
Sediment Quality Guidelines for the Protection of Aquatic Life		486 µg/kg dw PEL 170 µg/kg dw TEL	50-586 µg/kg dw	CCME (1999 and updates); ITT (2015)
Soil Quality Guidelines for the Protection of Environmental and Human Health				CCME (1999 and updates)
Agricultural, Residential and Parkland Land Uses		6600 µg/kg dw		
Commercial Land Use		24000 µg/kg dw		
Industrial Land Use		50000 µg/kg dw		
Air	0.3 µg/m ³ *			US EPA (1995)
Whole Body Lake Trout for the Protection of Aquatic Life		500 ng/g	121-233 ng/g	1987 GLWQA (IJC, 1987); McGoldrick and Murphy (2016)
Fish Fillet Human Health Criterion	0.3 µg/g ww	0.5 µg/g ww		(Evers et al., 2011); ITT (2015)
Blood: sensitive populations	5.8 µg/L	8 µg/L		ITT (2015)
Methylmercury				
Water Quality Guidelines for the Protection of Aquatic Life (freshwater)	1.4 µg/L CMC 0.77 µg/L CCC	0.004 µg/L		US EPA (1996); CCME (1999 and updates)
Tissue Residue Quality Guideline for the Protection of Wildlife Consumers of Aquatic Biota		33 µg/kg diet ww		CCME (1999 and updates)
US FDA Action Level	1 mg/kg			ITT (2015)

CCC = Criterion Continuous Concentration (chronic); CMC = Criterion Maximum Concentration (acute); dw = dry weight; MCL = Maximum Containment Level; PEL = Probable Effects Level; TEL = Threshold Effects Level; wt = wet weight

* this value is an inhalation reference concentration, not a standard, and has no regulatory standing. It is an estimate (with uncertainty spanning perhaps an order of magnitude) of a continuous inhalation exposure to the human population (including sensitive subgroups) that is likely to be without an appreciable risk of deleterious effects during a G-6 lifetime. It can be derived from a NOAEL, LOAEL, or benchmark concentration, with uncertainty factors generally applied to reflect limitations of the data used. Generally used in EPA's noncancer health assessments.

(https://cfpub.epa.gov/ncea/iris/iris_documents/documents/subst/0370_summary.pdf)

9 References

- AFN (2013). First Nations Biomonitoring Initiative National Results (2011). (http://www.afn.ca/uploads/files/afn_fnbi_en_-_2013-06-26.pdf). Ottawa, Ontario: Assembly of First Nations.
- AMAP (2011). Arctic Pollution 2011 (Mercury). (<http://hdl.handle.net/11374/634>). Arctic Monitoring and Assessment Programme
- ASTDR (1999). Toxicological Profile for Mercury. (<https://www.atsdr.cdc.gov/ToxProfiles/tp46.pdf>). Atlanta, GA: Agency for Toxic Substances and Disease Registry. U.S. Department of Health and Human Services, Public Health Service.
- ATSDR (2013). Addendum to the Toxicological Profile for Mercury (Alkyl and Dialkyl Compounds): Supplement to the 1999 Toxicological Profile for Mercury. (https://www.atsdr.cdc.gov/toxprofiles/mercury_organic_addendum.pdf). Atlanta, GA: Agency for Toxic Substances and Disease Registry. Division of Toxicology and Human Health Sciences.
- Basu, N. and J. Head (2010). "Mammalian wildlife as complementary models in environmental neurotoxicology." *Neurotoxicology and Teratology* 32(1): 114-119.
- Bender, M. and D. Narvaez (2016). Update on Global Mercury Production and Trade Trends, and Need for Improved Reporting. UNEP Supply/Storage Partnership Meeting, Madrid, Spain.
- Benoit, N., A. Dove, D. Burniston and D. Boyd (2016). "Tracking PCB Contamination in Ontario Great Lakes Tributaries: Development of Methodologies and Lessons Learned for Watershed Based Investigations." *Journal of Environmental Protection* 7(3).
- Berndt, M.E. (2003). *Mercury and Mining in Minnesota*. Minerals Coordinating Committee Final Report. Minnesota Department of Natural Resources Division of Lands and Minerals.
- Bhavsar, S.P., S.B. Gewurtz, D. J. McGoldrick, M.J. Keir, and S.M. Backus (2010). Changes in Mercury Levels in Great Lakes Fish Between 1970s and 2007. *Environ. Sci. Technol.*, 44, 3273–3279
- Carpenter, C., L. O'Connor, J. Elmer and D. DePinho (2011). Assessing the Impacts of the Mercury Export Ban Act of 2008 on the U.S. Mercury Recycling Industry. WM11 Global Achievements and Challenges in Waste Management, Phoenix, AZ.
- Carlson D. and D. Swackhamer. 2006. Results from the U.S. Great Lakes Fish Monitoring Program and effects of lake processes on bioaccumulative contaminant concentrations. *Journal of Great Lakes Research* 32: 370 – 385.
- CCME (2000). Canada-Wide Standards for Mercury Emissions. (http://www.ccme.ca/files/Resources/air/mercury/mercury_emis_std_e1.pdf). Quebec City, Quebec: Canadian Council of Ministers of the Environment. Government of Canada.
- CCME (2001). Canada-Wide Standard for Mercury-Containing Lamps. (http://www.ccme.ca/files/Resources/air/mercury/merc_lamp_standard_e.pdf). Winnipeg, Manitoba: Canadian Council of Ministers of the Environment. Government of Canada.
- CCME (2006). Canada-Wide Standards for Mercury Emissions from Coal-Fired Electric Power Generation Plants. (<http://www2.gnb.ca/content/dam/gnb/Departments/env/pdf/LandWaste-TerreDechets/Mercury-Mercure/CoalFiredElectricPowerGenerationPlants.pdf>). Canadian Council of Ministers of the Environment. Government of Canada.
- CCME (2007). Canada-Wide Standards for Mercury: A Report on Compliance and Evaluation Mercury from Dental Amalgam Waste, A Report of Progress Mercury Emissions and Mercury-Containing Lamps. (http://www.ccme.ca/files/Resources/air/mercury/2007_joint_hg_rpt_1.0_e.pdf). Canadian Council of Ministers of the Environment. Government of Canada.
- CCME (2010). Canada-Wide Standard for Mercury Emissions (Incineration & Base Metal Smelting)- 2010 Progress Report. (http://www.ccme.ca/files/Resources/air/mercury/mercury_incin_bms_2010_progress_rpt_e.pdf). Canadian Council of Ministers of the Environment. Government of Canada.

- CCME (1999 and updates). *Canadian Environmental Quality Guidelines*. Canadian Council of Ministers of the Environment. Available from: <http://cegg-rcqe.ccme.ca/en/index.html> (accessed June 2017).
- CCME (2016). Canada-Wide Standards for Mercury Emissions from Coal-Fired Electric Power Generation Plants: 2013/14 Progress Report (PN 1563; ISBN 978-1-77202-036-6). Canadian Council of Ministers of the Environment. Government of Canada.
- CEC (2015). Close-out Report: North American Regional Action Plan on Environmental Monitoring and Assessment, Chemicals Inventory and Mercury Activities in Mexico. (cec.org). Commission for Environmental Cooperation.
- Christensen, K. Y., B. A. Thompson, M. Werner, K. Malecki, P. Imm and H. A. Anderson (2015). "Levels of Nutrients in Relation to Fish Consumption Among Older Male Anglers in Wisconsin." *Environmental Research* 142: 542-548.
- Christensen, K. Y., B. A. Thompson, M. Werner, K. Malecki, P. Imm and H. A. Anderson (2016). "Levels of Persistent Contaminants in Relation to Fish Consumption Among Older Male Anglers in Wisconsin." *International Journal of Hygiene and Environmental Health* 219(2): 184-194.
- Cohen, M. D., R. S. Artz and R. R. Draxler (2007). Report to Congress: Mercury Contamination in the Great Lakes. (https://www.arl.noaa.gov/documents/reports/NOAA_GL_Hg.pdf). Silver Spring, MD: Air Resources Laboratory. National Oceanic and Atmospheric Administration, Office of Oceanic and Atmospheric Research.
- Cole, A., A. Steffen, C. Eckley, J. Narayan, M. Pilote, R. Tordon, J. Graydon, V. St. Louis, X. Xu and B. Branfireun (2014). "A Survey of Mercury in Air and Precipitation across Canada: Patterns and Trends." *Atmosphere* 5(3): 635.
- ECCC (2007). "Mercury Releases from Mercury Switches in End-of-Life Vehicles (Notice Completed)." *Canada Gazette, Part I* 141(52): 3556.
- ECCC (2010). "Notice Requiring the Preparation and Implementation of Pollution Prevention Plans in Respect of Mercury Releases from Dental Amalgam Waste." *Canada Gazette, Part I* 144(19).
- ECCC and Health Canada (2010). *Risk Management Strategy for Mercury - Highlights*. Government of Canada. Available from: https://www.ec.gc.ca/doc/mercure-mercury/1241/index_e.htm.
- ECCC (2014). Products Containing Mercury Regulations (SOR/2014-254). (SOR-2014-254). Environment and Climate Change Canada. Government of Canada.
- ECCC (2015). *Risk Management Strategy for Mercury-Containing Products*. Available from: <https://www.ec.gc.ca/lcpe-cepa/default.asp?lang=En&n=C54B4FE5-1&offset=1> (accessed September 2017).
- ECCC (2016a). Canadian Mercury Science Assessment - Executive Summary. (<https://www.ec.gc.ca/mercure-mercury/default.asp?lang=En&n=A2D7E54F-1>). Gatineau, Quebec: Environment and Climate Change Canada.
- ECCC (2016b). Canadian Mercury Science Assessment - Summary of Key Results. (<https://www.ec.gc.ca/mercure-mercury/default.asp?lang=En&n=32909A5D-1>). Gatineau, Quebec: Environment and Climate Change Canada.
- ECCC (2016c). *National Pollutant Release Inventory*. Environment and Climate Change Canada. Available from: <https://www.ec.gc.ca/inrp-npri/> (accessed September 2017).
- ECCC (2017a). *The National Atmospheric Chemistry Database and Analysis Facility*. Science and Technology Branch of Environment and Climate Change Canada. Available from: <https://www.ec.gc.ca/natchem/> (accessed September 2017).
- ECCC (2017b). "Notice of the Code of Practice for the Environmentally Sound Management of End-of-life Lamps Containing Mercury" *Canada Gazette Part I* 151(6).
- ECCC (2018). Products Containing Mercury Regulations: proposed amendments.

- (<https://www.canada.ca/en/environment-climate-change/services/canadian-environmental-protection-act-registry/products-mercury-regulations-proposed-amendments.html>) (accessed February 2018).
- ECCC (2018a). Air Pollutant Emission Inventory Report, 1996-2016. Available from: <https://www.canada.ca/content/dam/eccc/images/apei/apei-2018-en.pdf>
- ECCC and US EPA (2011). State of the Great Lakes 2011. (EPA 950-R-13-002). Cat No. En161-3/1-2011E-PDF. State of the Lakes Ecosystem Conferences (SOLEC).
- ECCC and US EPA (2017). State of the Great Lakes Highlights Report 2017. (https://binational.net/wp-content/uploads/2017/06/SOGL_17-EN.pdf). Cat. No. : En161-3E-PDF. State of the Lakes Ecosystem Conferences (SOLEC).
- ECE (2014). 1998 Protocol on heavy Metals, as amended on 13 December 2012. (ECE/EB.AIR/115). Executive Body for the Convention on Long-range Transboundary Air Pollution. United Nations Economic and Social Council.
- European Commission (2012). The International Negotiations on a Global Legally Binding Instrument on Mercury. (<http://ec.europa.eu/environment/chemicals/mercury/pdf/Report-Support-Hg-Negotiations.pdf>). Directorate-General Environment.
- Evers, D. C., J. G. Wiener, C. T. Driscoll, D. A. Gay, N. Basu, B. A. Monson, K. F. Lambert, H. A. Morrison, J. T. Morgan, K. A. Williams and A. G. Soehl (2011). Great Lakes Mercury Connections: The Extent and Effects of Mercury Pollution in the Great Lakes Region. (http://harvardforest.fas.harvard.edu/sites/harvardforest.fas.harvard.edu/files/publications/pdf/s/GLMC_FinalReport.pdf). Gorham, Maine: B. R. Institute.
- Federal Minister of the Environment and Canadian Dental Association (2002). Memorandum of Understanding Respecting the Implementation of the Canada-Wide Standard on Mercury for Dental Amalgam Waste. (https://www.ec.gc.ca/mercure-mercury/5910BAFF-FA15-40F9-B680-7195AD689A4D/EC_CDA_MOU.pdf). Ottawa, Canada: Minister of the Environment. Government of Canada.
- Ghandi, N., S.P. Bhavsar, R.W.K. Tang and G.B. Arhonditsis (2015). Projecting Fish Mercury Levels in the Province of Ontario, Canada and the Implications for Fish and Human Health. *Environ. Sci. Technol.* 49, 14494–14502
- Ginsberg, G. and B.F. Toal (2009). Quantitative Approach for Incorporating Methylmercury Risks and Omega-3 Fatty Acid Benefits in Developing Species-Specific Fish Consumption Advice. *Environ Health Perspect* 117:267–275. doi:10.1289/ehp.11368
- Ginsberg, G., (2016). Risk/Benefit Analysis of Great Lakes Fish for Neurodevelopmental Outcomes. Great Lakes Consortium for Fish Consumption Advisories. Minnesota Department of Health.
- GLRC (2010). Great Lakes Mercury Emission Reduction Strategy. (<http://www.glrppr.org/glmst/Mercury-Emissions-Reduction-Strategy.pdf>). Great Lakes Regional Collaboration.
- GLRC (2014). Great Lakes Mercury Emission Reduction Strategy Progress Report. (Document is not publicly available, but can be obtained upon request.). Great Lakes Regional Collaboration.
- Government of Canada (2016). *Chemicals Management Plan*. Available from: <https://www.canada.ca/en/health-canada/services/chemical-substances/chemicals-management-plan.html> (accessed September 2017).
- Great Lakes Consortium for Fish Consumption Advisories 2018. <http://www.health.state.mn.us/divs/eh/fish/consortium/members.html>
- Great Lakes Interagency Task Force (2015). Great Lakes Restoration Initiative Report to Congress and the President- Fiscal Years 2010-2014. (https://www.glri.us/pdfs/21050720-report_to_congress.pdf). G. L. R. Initiative.
- Health Canada (1986). Guidelines for Canadian Drinking Water Quality: Guideline Technical Document – Mercury. (<https://www.canada.ca/en/health-canada/services/publications/healthy->

- [living/guidelines-canadian-drinking-water-quality-guideline-technical-document-mercury.html#a1](#)). Government of Canada.
- Huang, J., S.N. Lyman, J. Stamenkovic Hartman and M. Sexauer Gustin (2014). A review of passive sampling systems for ambient air mercury measurements *Environ. Sci.: Processes Impacts*, 2014, 16, 374. DOI: 10.1039/c3em00501a
- IJC (1987). Revised Great Lakes Water Quality Agreement of 1978. (http://www.ijc.org/files/tinymce/uploaded/GLWQA_e.pdf). Ottawa: International Joint Commission. United States and Canada.
- International Joint Commission (2015). Atmospheric Deposition of Mercury in the Great Lakes. (http://publications.gc.ca/collections/collection_2016/cmi-ijc/E95-2-21-2015-eng.pdf).
- ITT (2015). Binational Summary Report: Mercury. (<https://binational.net/wp-content/uploads/2015/05/EN-Mercury-Binational-Summary-Report-Final-Draft.pdf>). Identification Task Team.
- Jeremiason, J. (2017). A Synthesis of Mercury Trends in Lake Superior. (Document is not publicly available, but can be obtained upon request.).
- Kim, S., P. A. Thiessen, E. E. Bolton, J. Chen, G. Fu, A. Gindulyte, L. Han, J. He, S. He, B. A. Shoemaker, J. Wang, B. Yu, J. Zhang and S. H. Bryant (2016). "PubChem Substance and Compound databases." *Nucleic Acids Res* 44(D1): D1202-1213.
- Lake Superior Partnership (2016). Lake Superior Lakewide Action and Management Plan 2015-2019. (Cat. No. En164-52/2016E-PDF).
- Lamborg, C.H., C.R. Hammerschmidt, K.L. Bowman, G.J. Swarr, K.M. Munson, D.C. Ohnemus, P.J. Lam, L.E. Heimbürger, M.J. A. Rijkenberg and M.A. Saito (2014). A global ocean inventory of anthropogenic mercury based on water column measurements. *Nature* volume 512, pages 65–68. doi:10.1038/nature13563
- Lepak, R.F., R. Yin, D.P. Krabbenhoft, J.M. Ogorek, J.F. DeWild, T.M. Holsen, and J.P. Hurley (2015). Use of Stable Isotope Signatures to Determine Mercury Sources in the Great Lakes. *Environmental Science & Technology Letters* 2 (12), 335-341
DOI: 10.1021/acs.estlett.5b00277
- Mahaffey, K. R. (2005). "Mercury Exposure: Medical and Public Health Issues." *Transactions of the American Clinical and Climatological Association* 116: 127-154.
- McGoldrick, D. J. and E. W. Murphy (2016). "Concentration and Distribution of Contaminants in Lake Trout and Walleye from the Laurentian Great Lakes (2008–2012)." *Environmental Pollution* 217: 85-96.
- McLagan, D.S., M.E.E. Mazur, C.P.J. Mitchell, and F. Wania (2016). Passive air sampling of gaseous elemental mercury: a critical review. *Atmos. Chem. Phys.*, 16, 3061–3076, doi:10.5194/acp-16-3061-2016
- Menyas, P. (2016). Canada Proposes Restrictions on Mercury Exports. *Bloomberg BNA: Daily Environment Report*, Bloomberg BNA.
- Michigan DEQ and US EPA (2013). Statewide Michigan Mercury TMDL. (http://www.michigan.gov/documents/deq/wrd-swms-hgtmdl-draft_415360_7.pdf). Michigan Department of Environmental Quality. US Environmental Protection Agency Region 5.
- Monson, B. A., D. F. Staples, S. P. Bhavsar, T. M. Holsen, C. S. Schrank, S. K. Moses, D. J. McGoldrick, S. M. Backus and K. A. Williams (2011). "Spatiotemporal trends of mercury in walleye and largemouth bass from the Laurentian Great Lakes Region." *Ecotoxicology* 20(7): 1555-1567.
- NADP (2011). Monitoring Mercury Deposition- A Key Tool to Understanding the Link between Emissions and Effects. (IEM 2005-03 and NADP Brochure 2005-01). National Atmospheric Deposition Program. NEI (2011). US EPA's National Emissions Inventory 2011 Report. Available from: <https://www.epa.gov/air-emissions-inventories/2011-national-emissions-inventory-nei-data>

- OMOECC (2013). Guide to Eating Ontario Sport Fish, 2013-2014. (www.ontario.ca/fishguide). Ontario Ministry of Environment and Climate Change. Government of Canada.
- OMOECC (2018). <https://www.ontario.ca/environment-and-energy/eating-ontario-fish>.
- Park, J.-D. and W. Zheng (2012). "Human Exposure and Health Effects of Inorganic and Elemental Mercury." *Journal of Preventive Medicine and Public Health* 45(6): 344-352.
- Quicksilver Caucus (2012). Thrid Compendium of States' Mercury Activities. (http://www.michigan.gov/documents/deq/Third_Compndium_of_States_Mercury_Activities_FINAL_Oct_2012_407789_7.pdf). Washington D.C.: The Environmental Council of the States.
- Raymond, M., K. Christensen, B. Thompson and H. Anderson (2017). "Changes in Hair Mercury Levels Among Women of Child-Bearing Age Following an Educational Intervention." *Journal of Occupational and Environmental Medicine* 59(6): 528-534.
- Schroeder, W. H. and J. Munthe (1998). "Atmospheric mercury—An overview." *Atmospheric Environment* 32(5): 809-822.
- Sibbald, B. (2016). "Doctors call to phase out coal-fired electricity." *Canadian Medical Association Journal* 188(17-18): E423-E424.
- Statistics Canada (2016). *Canadian Health Measures Survey (CHMS)*. Available from: <http://www23.statcan.gc.ca/imdb/p2SV.pl?Function=getSurvey&Id=148760> (accessed August 2017).
- Turyk, M. E., S. P. Bhavsar, W. Bowerman, E. Boysen, M. Clark, M. Diamond, D. Mergler, P. Pantazopoulos, S. Schantz and D. O. Carpenter (2012). "Risks and Benefits of Consumption of Great Lakes Fish." *Environmental Health Perspectives* 120(1): 11-18.
- UN Environment (2017a). *Minamata Convention on Mercury*. Available from: <http://www.mercuryconvention.org/> (accessed June 2017).
- UN Environment (2017b). Toolkit for Identification and Quantification of Mercury Releases: Guideline for Inventory Level 1. (<https://wedocs.unep.org/bitstream/handle/20.500.11822/14777/Hg-Toolkit-Guideline-IL1-January2017.pdf?sequence=1&isAllowed=y>). Geneva, Switzerland: UN Environment Chemicals Branch.
- UN Environment, (2017c). Global mercury supply, trade and demand. United Nations Environment Programme, Chemicals and Health Branch. Geneva, Switzerland.
- UNECE (2017). *Protocol on Heavy Metals- The 1998 Aarhus Protocol on Heavy Metals*. Available from: http://www.unece.org/env/lrtap/hm_h1.html (accessed September 2017).
- US Ecology (2017). *ELVS Mercury Switch Program*. Available from: <https://www.usecology.com/Services/Environmental-Services/Recycling/ELVS-Mercury-Switch-Program.aspx> (accessed
- US EPA (1995). Integrated Risk Information System (IRIS), Chemical Assessment Summary: Mercury, elemental; CASRN 7439-97-6. Available at: https://cfpub.epa.gov/ncea/iris/iris_documents/documents/subst/0370_summary.pdf
- US EPA (1996). Water Quality Criteria Documents for the Protection of Aquatic Life in Ambient Wawter. (EPA-820-B-96-001). Office of Water. US Environmental Protection Agency.
- US EPA (2004). Results of the Lake Michigan Mass Balance Study: Biphenyls and Trans-Nonachlor Data Report. (EPA 905 R-01-011). Chicago, IL: Great Lakes National Program Office. US Environmental Protection Agency.
- US EPA (2007). *Background Paper for Stakeholder Panel to Address Options for Managing U.S. Non-Federal Supplies of Commodity-Grade Mercury*. Office of Prevention, Pesticides and Toxic Substances .
- US EPA (2009). Report to Congress: Potential Export of Mercury Compounds from the United States for Conversion to Elemental Mercury. (<https://www.epa.gov/mercury/2009-epa-report-congress->

- [potential-export-mercury-compounds-united-states-conversion](#)). Washington, D.C.: Office of Pollution Prevention and Toxic Substances. US Environmental Protection Agency.
- US EPA (2010). 2010-2014 Pollution Prevention (P2) Program Strategic Plan. (<https://www.epa.gov/sites/production/files/documents/p2strategicplan2010-14.pdf>). US Environmental Protection Agency.
- US EPA (2014a). Data from the Great Lakes Fish Monitoring and Surveillance Program: Mercury.
- US EPA (2014b). *Toxics Release Inventory (TRI) Program*. Available from: <https://www.epa.gov/toxics-release-inventory-tri-program> (accessed June 2016).
- US EPA (2015a). Find Out What's Happening in Your Neighborhood Using EPA's Toxics Release Inventory (TRI). (https://www.epa.gov/sites/production/files/2015-10/documents/2015_tri_for_communities_fact_sheet_final.pdf). US Environmental Protection Agency.
- US EPA (2015b). Lake Michigan Lakewide Action and Management Plan Annual Report 2015. (<https://binational.net>).
- US EPA (2015c). List of Lists: Consolidated List of Chemicals Subject to the Emergency Planning and Community Right-To-Know Act (EPCRA), Comprehensive Environmental Response, Compensation and Liability Act (CERCLA) and Section 112(r) of the Clean Air Act. (EPA 550-B-15-001). Washington, D.C.: Office of Solid Waste and Emergency Response. US Environmental Protection Agency.
- US EPA (2016a). *Getting Work Done at AOCs: How are the Federal GLRI Agencies Implementing the AOC Program?* 2016 Great Lakes AOCs Conference, Dearborn, MI, US Environmental Protection Agency, Great Lakes National Program Office.
- US EPA (2016b). *Great Lakes Environmental Database (GLENDa)*. Available from: <https://www.epa.gov/great-lakes-legacy-act/great-lakes-environmental-database-glenda> (accessed June 2016).
- US EPA (2016c). *Great Lakes Monitoring: The Integrated Atmospheric Deposition Network (IADN)*. Available from: <https://www3.epa.gov/greatlakes/monitoring/air2/index.html> (accessed June 2016).
- US EPA (2016d). *Lakewide Action and Management Plans*. Available from: <https://www.epa.gov/greatlakes/lakewide-action-and-management-plans> (accessed August 2016).
- US EPA (2017a). Effluent Limitation Guidelines and Standards for the Dental Category; 40 CFR 441. (EPA-HQ-OW-2014-0693; FRL-9957-10-OW). US Environmental Protection Agency.
- US EPA (2017b). *Mercury in Your Environment*. United States Environmental Protection Agency. Available from: <https://www.epa.gov/mercury> (accessed June 21).
- US EPA (2017c). Mercury U.S. Inventory Report: Supply, Use, and Trade 2017. Office of Chemical Safety and Pollution Prevention. <https://www.regulations.gov/document?D=EPA-HQ-OPPT-2017-0127-0002>
- US EPA (2017d). *Mercury and Air Toxics Standards (MATS)*. Available from: <https://www.epa.gov/mats> (accessed September 2017).
- US EPA (2018). 2014 National Emissions Inventory (NEI) version 2 Technical Support Document (TSD).
- US EPA and Canada (2009). Great Lakes Binational Toxics Strategy- 2009 Biennial Report. (EN 161-1/2-2009E-PDF; 978-1-100-17486-0). US Environmental Protection Agency, and the Government of Canada.
- US EPA and ECCC (1997). The Great Lakes Binational Toxics Strategy. (<https://archive.epa.gov/greatlakes/p2/web/pdf/bnssign.pdf>).

- US FDA (2014). A Quantitative Assessment Of The Net Effects On Fetal Neurodevelopment From Eating Commercial Fish (As Measured by IQ and also by Early Age Verbal Development in Children). <https://www.fda.gov/Food/FoodborneIllnessContaminants/Metals/ucm393211.htm>
- Zananski, T.J., T. M. Holsen, P.K. Hopke and B. S. Crimmins (2011). Mercury temporal trends in top predator fish of the Laurentian Great Lakes. *Ecotoxicology* 20:1568–1576 DOI 10.1007/s10646-011-0751-9 *Ecotoxicology* (2011) 20:1568–1576 DOI 10.1007/s10646-011-0751-9
- Zhang, Y., D.J. Jacob, H.M. Horowitz, L. Chen, H.M. Amos, D.P. Krabbenhoft, F. Slemr, V.L. St. Louis and E.M. Sunderland (2016). Observed decrease in atmospheric mercury explained by global decline in anthropogenic emissions. *Proceedings of the National Academy of Sciences*, January, 113 (3) 526-531. <https://doi.org/10.1073/pnas.1516312113>
- Zhou, C., M.D. Cohen, B.A. Crimmins, H. Zhou, T.A. Johnson, P.K. Hopke, and T.M. Holsen (2017). Mercury Temporal Trends in Top Predator Fish of the Laurentian Great Lakes from 2004 to 2015: Are Concentrations Still Decreasing? *Environmental Science & Technology* 2017 51 (13), 7386-7394 DOI: 10.1021/acs.est.7b00982